

Datasheet: AAM64B BATCH NUMBER 170601

Description:	GOAT ANTI MOUSE SDF-1 ALPHA:Biotin
Specificity:	SDF-1 ALPHA
Other names:	CXCL12a
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	•			0.25 - 1.0ug/ml
Immunoprecipitation				
Western Blotting	•			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

G conjugated to Biotin - lyophilised
te with 0.5ml sterile PBS containing 0.1% bovine serum albumin d be taken during reconstitution as the protein may appear as a film at the he vial. Bio-Rad recommend that the vial is gently mixed after reconstitutior

Antiserum Preparation Antiserum to mouse SDF-1 alpha were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Preservative Stabilisers None present Carrier Free Yes Approx. Protein Concentrations IgG concentration 0.1 mg/ml after reconstitution Immunogen Recombinant mouse SDF-1 alpha expressed in E. coli (PMP77). External Database Links UniProt:	Buffer Solution	Phosphate buffered saline
Approx. Protein Concentrations IlgG concentration 0.1mg/ml after reconstitution Recombinant mouse SDF-1 alpha expressed in E. coli (PMP77). External Database Links UniProt: P40224 Related reagents Entrez Gene: 20315 Cxcl12 Related reagents Synonyms Sdf1 RRID AB_2088307 Specificity Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein-coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162		None present
Immunogen Recombinant mouse SDF-1 alpha expressed in E. coli (PMP77). External Database Links UniProt: P40224 Related reagents Entrez Gene: 20315 Cxcl12 Related reagents Synonyms Sdf1 RRID AB_2088307 Specificity Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein—coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	Carrier Free	Yes
External Database Links UniProt: P40224 Related reagents Entrez Gene: 20315 Cxcl12 Related reagents Synonyms Sdf1 RRID AB_2088307 Specificity Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein—coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162		IgG concentration 0.1mg/ml after reconstitution
Links P40224 Related reagents	Immunogen	Recombinant mouse SDF-1 alpha expressed in E. coli (PMP77).
Specificity Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein—coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162		P40224 Related reagents Entrez Gene:
Specificity Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein—coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	Synonyms	Sdf1
known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein—coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	RRID	AB_2088307
ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	Specificity	known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons.
ELISA with AAM64 as the capture antibody and PMP77 as the standard. Storage Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162		as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells.
After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	ELISA	
recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	Storage	
Health And Safety Information Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AAM64B 10162		recommended. Avoid repeated freezing and thawing as this may denature the antibody.
Information https://www.bio-rad-antibodies.com/SDS/AAM64B 10162	Guarantee	12 months from date of despatch
Regulatory For research purposes only	-	https://www.bio-rad-antibodies.com/SDS/AAM64B

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 America

Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.comd a Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com

То

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372430:200706'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint