

Datasheet: AHP031F BATCH NUMBER 155035

Description:	SHEEP ANTI HUMAN C3c:FITC
Specificity:	C3c
Other names:	COMPLEMENT COMPONENT 3c
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

Product Details

Applications	This product has been derived from testing wi communications from t information. For genera	thin our labora he originators	atories, p . Please	eer-reviewed public refer to references	indicated for further
	rad-antibodies.com/pro	tocols.			
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				
	Immunohistology - Frozer	n •			1/50 - 1/100
	Immunohistology - Paraff	in			
	Where this antibody han necessarily exclude its a guide only. It is recorn system using appropria	use in such p nmended that	orocedure the user	es. Suggested worki titrates the antibod	ing dilutions are given as
Target Species	Human				
Product Form	Purified IgG conjugated	d to Fluoresce	ein Isothio	ocyanate Isomer 1 (FITC) - liquid
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)	
	FITC	490		525	-
Antiserum Preparatio	n Antisera to human C3c purified antigen. Purifie		• •		
Buffer Solution	Phosphate buffered sa	line			
Preservative Stabilisers	0.099% Sodium Azide				
Approx. Protein	IgG concentration 10 n	ng/ml			

Concentrations

Immunogen	Human C3 purified from serum.
External Database Links	UniProt:P01024Related reagentsEntrez Gene:718C3Related reagents
Synonyms	CPAMD1
RRID	AB_321213
Specificity	Sheep anti Human C3c antibody recognizes the C3c component of human complement, formed as a result of the inactivation of C3b. Sheep anti Human C3c antibody may be used for the detection of C3 deposits in tissues following complement activation.
Histology Positive Control Tissue	Human skin
References	 Kennedy, M.W. & Kuo, Y.M. (1988) The surfaces of the parasitic nematodes <i>Trichinella spiralis</i> and <i>Toxocara canis</i> differ in the binding of post-C3 components of human complement by the alternative pathway. <u>Parasite Immunol. 10:459-63.</u> Oyeyinka, G.O. <i>et al.</i> (2003) The effects of ageing on the immune response to <i>Schistosoma haematobium</i> and hookworm by measuring circulating immune complexes, C3, IgG, IgA and IgM levels in residents of Omi dam area of Kogi State, Nigeria. <u>Afr J Med Med Sci. 32: 263-7.</u> Rojana-Udomsart, A. <i>et al.</i> (2013) Complement-mediated muscle cell lysis: A possible mechanism of myonecrosis in anti-SRP associated necrotizing myopathy (ASANM). <u>J Neuroimmunol. 264: 65-70.</u> Akinlade, K.S. <i>et al.</i> (2004) Circulating immune complexes, immunoglobulin classes (IgG, IgA and IgM) and complement components (C3c, C4 and Factor B) in diabetic Nigerians. <u>West Afr J Med. 23 (3): 253-5.</u> Rojana-udomsart, A. <i>et al.</i> (2013) Complement-mediated muscle cell lysis: a possible mechanism of myonecrosis in anti-SRP associated necrotizing myopathy (ASANM). J <u>Neuroimmunol. 264 (1-2): 65-70.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

urposes only		
	•	9 (0) 89 8090 95 21 9 (0) 89 8090 95 50
		antibody_sales_de@bio-rad.cor
r	Fax: +44 (0)1865 852 739 m Email: antibody_sales_uk@bio-ra	Fax: +44 (0)1865 852 739 Fax: +4 n Email: antibody_sales_uk@bio-rad.com Email: a his product, please use our online search tool at: bio-rad-a

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint