

Datasheet: AHP1003

Description:	GOAT ANTI HUMAN LNK (N-TERMINAL)
Specificity:	LNK (N-TERMINAL)
Other names:	LYMPHOCYTE ADAPTOR PROTEIN
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			0.3 - 1.0ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Antisera to LNK (NT) were raised by repeated immunisation of Goat with highly purified antigen. Purified IgG prepared by affinity chromatography.
Buffer Solution	TRIS buffered saline
Preservative	0.02% Sodium Azide
Stabilisers	0.5% Bovine Serum Albumin
Approx. Protein	IgG concentration 0.5mg/ml

Concentrations

Immunogen Peptide sequence NGPALQPSSPC from LNK amino-terminus.

External Database Links

UniProt:

[Q9UQQ2](#) [Related reagents](#)

Entrez Gene:

[10019](#) SH2B3 [Related reagents](#)

Synonyms LNK

RRID AB_2270411

Specificity

Goat anti Human LNK antibody recognizes human SH2B adapter protein 3, also known as Lymphocyte-specific adapter protein Lnk or Lymphocyte adapter protein. LNK is a 575 amino acid SH2B adapter protein containing a single PH domain and a single SH2 domain. Goat anti Human LNK antibody binds to an epitope within the N-terminal region of LNK, which regulates cytokine receptor-mediated and growth factor pathways, and plays an essential role in B cell lymphopoiesis.

LNK mRNA is preferentially expressed in lymph node and spleen lymphocytes and the expression of LNK in vascular endothelial cells (ECs), increases in response to TNF α . LNK acts as a negative regulator of the pro-inflammatory molecules VCAM-1 and E-selectin induced by TNF α , modulation of P13-kinase and MAPK ERK1/2 activity. LNK is also involved in the negative regulation of Thrombopoietin-mediated cell proliferation and endomitosis in hematopoietic cell lines.

Western Blotting

Goat anti Human LNK antibody, directed against the N-teend of the molecule, detects a band of approximately 70 kDa in human MOLT4 cell lysates. (Predicted Mwt. 63.2 kDa). A band of 68 kDa was resolved in HUVECs ([Fitau et al. 2006](#)).

Suitable antibodies for loading controls include :

Human anti Human actin β , [HCA147](#) or Rat anti tubulin α , [MCA77G](#).

References

1. Fitau, J. *et al.* (2005) The adaptor protein Lnk modulates endothelial cell activation. [Nephrol Ther. 1: 228-33.](#)
2. Wan, M. *et al.* (2006) TNF- α Induces Lnk Expression Through PI3K-Dependent Signaling Pathway in Human Umbilical Vein Endothelial Cells [J Surg Res. 136: 53-7.](#)
3. Fitau, J. *et al.* (2006) The adaptor molecule Lnk negatively regulates tumor necrosis factor-alpha-dependent VCAM-1 expression in endothelial cells through inhibition of the ERK1 and -2 pathways. [J Biol Chem. 281 \(29\): 20148-59.](#)
4. Wan, M. *et al.* (2007) Eicosapentaenoic acid inhibits TNF-alpha-induced Lnk expression in human umbilical vein endothelial cells: involvement of the PI3K/Akt pathway. [J Nutr Biochem. 18: 17-22.](#)
5. Gery, S. *et al.* (2007) Adaptor protein Lnk negatively regulates the mutant MPL,

- MPLW515L associated with myeloproliferative disorders. [Blood. 110: 3360-4.](#)
6. Gery, S. *et al.* (2009) Lnk inhibits myeloproliferative disorder-associated JAK2 mutant, JAK2V617F. [J Leukoc Biol. 85: 957-65.](#)
7. Chatelais, M. *et al.* (2011) Gene transfer of the adaptor Lnk (SH2B3) prevents porcine endothelial cell activation and apoptosis: implication for xenograft's cytoprotection. [Xenotransplantation. 18: 108-20.](#)
8. Devallière, J. *et al.* (2012) LNK (SH2B3) is a key regulator of integrin signaling in endothelial cells and targets α -parvin to control cell adhesion and migration. [FASEB J. 26: 2592-606.](#)

Further Reading 1. Tong, W. & Lodish, H.F. (2004) Lnk inhibits Tpo-mpl signaling and Tpo-mediated megakaryocytopoiesis. [J Exp Med. 200 \(5\): 569-80.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10058 available at: <https://www.bio-rad-antibodies.com/SDS/AHP1003>
10058

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M428128:240301'

Printed on 01 Mar 2024