

Datasheet: AHP1031B BATCH NUMBER 151173

Description:	RABBIT ANTI HUMAN G-CSF:Biotin
Specificity:	G-CSF
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen				
Immunohistology - Paraffin	•			
ELISA	•			0.25 - 1.0ug/ml
Immunoprecipitation			•	
Western Blotting	•			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form Purified IgG conjugated to Biotin - Iyophilised Reconstitution Reconstitute with 0.5ml sterile PBS containing 0.1% Bovine Serum Albumin. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution For long term storage the addition of 0.09% sodium azide is recommended. Antiserum Preparation Antisera to human G-CSF were raised by repeated immunisations of rabbits with high purified antigen. Purified IgG prepared by affinity chromatography.	Target Species	Human	
Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution For long term storage the addition of 0.09% sodium azide is recommended. Antiserum Preparation Antisera to human G-CSF were raised by repeated immunisations of rabbits with high	Product Form	Purified IgG conjugated to Biotin - lyophilised	
· ·	Reconstitution	Care should be taken during reconstitution as the protein may bottom of the vial. Bio-Rad recommend that the vial is gently n	appear as a film at the nixed after reconstitution.
	Antiserum Preparation	• •	• •
Buffer Solution Phosphate buffered saline	Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml after reconstitution.
Immunogen	Recombinant human G-CSF
External Database Links	UniProt: P09919 Related reagents Entrez Gene: 1440 CSF3 Related reagents
Synonyms	C17orf33, GCSF
RRID	AB_2083735
Specificity	Rabbit anti Human G-CSF antibody recognizes human G-CSF (Granulocyte colonystimulating factor), a 19.6kDa haematopoietic growth factor and member of the CSF family of hormone-like glycoproteins, which stimulates the development of bone marrow progenitor cells and in particular those committed to neutrophil lineage. The ability of G-CSF to stimulate the proliferation/differentiation and function of neutrophil granulocyte progenitor cells and survival of mature neutrophils, has resulted in G-CSF becoming an important factor in patient treatment following allogeneic (bone marrow/stem cell) transplantation and chemotherapy. To yield one-half maximal inhibition [ND ₅₀] of the biological activity of hG-CSF (0.5 ng/ml), a concentration of 0.04-0.06 ug/ml of this antibody is required.
ELISA	This biotinylated human G-CSF antibody may be used in a direct ELISA or as the detection reagent in a sandwich ELISA with our <u>purified human G-CSF antibody</u> (AHP1031) as the capture reagent and <u>recombinant human G-CSF</u> (PHP082B) as the standard.
Further Reading	 Hill, C.P. et al. (1993) The structure of granulocyte-colony-stimulating factor and its relationship to other growth factors. Proc Natl Acad Sci U S A. 90 (11): 5167-71. Morton, J. et al. (2001) Granulocyte-colony-stimulating factor (G-CSF)-primed allogeneic bone marrow: significantly less graft-versus-host disease and comparable engraftment to G-CSF-mobilized peripheral blood stem cells. Blood. 98 (12): 3186-91.
Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C.
	This product should be stored undiluted. Storage in frost-free freezers is not

recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	6 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AHP1031B 10162
Regulatory	For research purposes only

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372460:200706'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint