

Datasheet: AHP1076 BATCH NUMBER 150303

Description:	GOAT ANTI STAT3 (C-TERMINAL)
Specificity:	STAT3 (C-TERMINAL)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been re derived from testing withi communications from the information. For general rad-antibodies.com/proto	in our labo e originato protocol re	oratories, ors. Pleas	peer-reviewed publica e refer to references in	tions or personal ndicated for further
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			-	
	Immunohistology - Frozen (1)	•			10ug/ml
	Immunohistology - Paraffin	•			
	ELISA	-			1/32,000
	Immunoprecipitation			•	
	Western Blotting	•			0 - 5ug/ml
	a guide only. It is recomn system using appropriate (1) Overnight staining is	e negative	/positive	controls.	for use in their own
Target Species	Human				
Species Cross Reactivity	Reacts with: Mouse Based on sequence simil N.B. Antibody reactivity a reactivity is derived from personal communications further information.	and workin testing wi	ng conditi thin our la	ons may vary between aboratories, peer-revie	wed publications or
Product Form	Purified IgG - liquid				

Antiserum Preparation Antisera to human STAT3 (CT) were raised by repeated immunisations of goats with

Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.02% Sodium Azide 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Peptide sequence DMELTSECARSPM corresponding to the C-terminal region of STAT3 (NP_003141.2; NP_644805.1).
External Database Links	UniProt: <u>P40763</u> <u>Related reagents</u> Entrez Gene: <u>6774</u> STAT3 <u>Related reagents</u>
Synonyms	APRF
RRID	AB_2196584
Specificity	Goat anti Mouse STAT3 antibody recognizes an epitope within the C-terminal (CT) region of signal transducer and activator of transcription 3 (STAT3), an 80-85 kDa member of the STAT protein family, which plays an important role in cell growth, survival differentiation and transformation, transducing signals from cytokine receptors to the nucleus.
	interleukin-5, interleukin-6, epidermal growth factor and leukaemia inhibitory factor. It is expressed in the heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.
Histology Positive Control Tissue	Mouse Thymus
Western Blotting	Goat anti STAT3 antibody detects a band of approximately 80-85kDa in A431 cell lysates. A minimum incubation time of 1 hour is recommended with this antibody.
Further Reading	1. Levy, D.E. & Lee, C.K. (2002) What does Stat3 do? <u>J Clin Invest. 109 (9): 1143-8.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we

Health A	nd Safety Mat	erial Safetv Datas	heet documentation #10)058 available	at:
Informati	-	•	ntibodies.com/SDS/AHF		
	100				
Regulato	ry For	research purpose	s only		
Relate	d Products				
	nended Second	ary Antibodies			
Recomn		-			
Recomn Rabbit Ar	nended Second	-		Europe	Tel: +49 (0) 89 8090 95 21
Recomn Rabbit Ar	n ended Second nti Goat IgG (Fc) (STAR122) <u>FITC</u>	, <u>HRP</u>	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint