

Datasheet: AHP1081

Description:	GOAT ANTI HUMAN MMP-7
Specificity:	MMP-7
Other names:	MATRILYSIN
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/64000
Immunoprecipitation			▪	
Western Blotting	▪			0.03 - 0.1ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG - liquid

Antiserum Preparation Antisera to human MMP7 were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution TRIS buffered saline

Preservative 0.02% Sodium Azide
Stabilisers 0.5% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.5 mg/ml

Immunogen Peptide sequence C-QKLYGKRSNSRKK corresponding to the C-terminus of MMP7 (NP_002414.1).

External Database Links **UniProt:**

[P09237](#) [Related reagents](#)

Entrez Gene:

[4316](#) MMP7 [Related reagents](#)

Synonyms MPSL1, PUMP1

RRID AB_2144333

Specificity **Goat anti Huma MMP7 antibody** recognises matrix metalloproteinase 7 (MMP7), a ~30 kDa member of the MMP family. MMP's are secreted as inactive zymogens which are activated when proteolytically cleaved by extracellular proteinases. They are involved in processes such as embryonic development, reproduction and tissue remodelling where they breakdown the extracellular matrix. This action is also exploited by diseases processes including arthritis and metastasis. MMP7 degrades casein, fibronectin, gelatin types I, III, IV, V, and activates procollagenase. This metalloproteinase has been identified as an activator of the Notch signalling pathway, and is also involved in wound healing and the regulation of defensins in intestinal mucosa.

Western Blotting AHP1081 detects a band of approximately 28kDa in human breast, ovary and placenta cell lysates. A minimum incubation time of 1 hour is recommended with this antibody.

Further Reading 1. Gaire, M. *et al.* (1994) Structure and expression of the human gene for the matrix metalloproteinase matrilysin. [J Biol Chem. 269 \(3\): 2032-40.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10058 available at:
10058: <https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M363804:200529'

Printed on 11 Aug 2020
