

Datasheet: AHP1100

Description:	GOAT ANTI HUMAN PAI-1
Specificity:	PAI-1
Other names:	SERPINE1
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			4.0 - 6.0 ug/ml
ELISA	▪			1/32000
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.3 ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This antibody requires heat-mediated antigen retrieval prior to staining paraffin sections. Citrate buffer pH6.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Based on sequence similarity, is expected to react with: Dog, Chimpanzee
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Antiserum Preparation

Antisera to human PAI-1 were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.02% Sodium Azide 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Peptide with sequence C-GFKIDDKGMAPALRH, corresponding to the internal region of PAI-1 (NP_000593.1).
External Database Links	<p>UniProt: P05121 Related reagents</p> <p>Entrez Gene: 5054 SERPINE1 Related reagents</p>
Synonyms	PAI1, PLANH1
RRID	AB_808706
Specificity	<p>Goat anti human PAI-1 antibody recognizes an internal epitope within human plasminogen activator inhibitor 1 (PAI-1), also known as serpin E1. PAI-1 is a 402 amino acid ~45 kDa serine protease inhibitor and the principle inhibitor of tissue plasminogen activator (tPA) and urokinase (uPA)</p> <p>PAI-1 is present in plasma, platelets and endothelial cells, is expressed at increased levels in various disease states (Kruithof et al. 1988), and has been linked to an increased occurrence of thrombosis in obesity and coronary artery disease (Oseroff et al. 1989). Defects in PAI-1 are characterized by abnormal bleeding (Takahashi et al. 1996).</p> <p>Goat anti human PAI-1 antibody has been used successfully for the detection of PAI-1 in human tissues by immunohistochemistry, western blotting and ELISA.</p>
Histology Positive Control Tissue	Human Placenta
Western Blotting	Goat anti human PAI-1 detects a band of approximately 45kDa in human duodenum, liver and lung lysates. A minimum incubation time of 1 hour is recommended with this antibody.
References	<ol style="list-style-type: none"> De Taeye, B. <i>et al.</i> (2005) Plasminogen activator inhibitor-1: a common denominator in obesity, diabetes and cardiovascular disease. Curr Opin Pharmacol. 5 (2): 149-54. Berg, D. <i>et al.</i> (2012) Profiling signalling pathways in formalin-fixed and paraffin-embedded breast cancer tissues reveals cross-talk between EGFR, HER2, HER3 and uPAR. J Cell Physiol. 227: 204-12. Wolff, C. <i>et al.</i> (2011) Signalling networks associated with urokinase-type plasminogen activator (uPA) and its inhibitor PAI-1 in breast cancer tissues: new insights from protein

microarray analysis. [J Pathol. 223: 54-63.](#)

4. Malinowsky, K. *et al.* (2014) uPA and PAI-1-Related Signaling Pathways Differ between Primary Breast Cancers and Lymph Node Metastases. [Transl Oncol. 5: 98-104.](#)

5. Berg, D. *et al.* (2011) Discovery of new molecular subtypes in oesophageal adenocarcinoma. [PLoS One. 6 \(9\): e23985.](#)

Further Reading 1. Lijnen, H.R. (2005) Pleiotropic functions of plasminogen activator inhibitor-1. [J Thromb Haemost. 3 \(1\): 35-45.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10058 available at: 10058: <https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[100x ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382785:210513'

Printed on 11 Jul 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)