

Datasheet: AHP1124

BATCH NUMBER 162929

Description:	GOAT ANTI SNAP-25 (C-TERMINAL)
Specificity:	SNAP-25 (C-TERMINAL)
Other names:	SYNAPTOSOMAL-ASSOCIATED PROTEIN
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			2.0 - 4.0ug/ml
ELISA	▪			1/32000
Immunoprecipitation			▪	
Western Blotting	▪			0.01 - 0.03ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Heat-mediated antigen retrieval is required prior to staining paraffin sections with this antibody. Bio-Rad recommend citrate buffer, pH6.0, for this purpose.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Mouse</p> <p>Based on sequence similarity, is expected to react with:Rat, Dog</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Antiserum Preparation Antisera to human SNAP-25 were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution TRIS buffered saline

Preservative 0.02% Sodium Azide
Stabilisers 0.5% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.5 mg/ml

Immunogen Peptide sequence C-DEANQRATKMLGSG corresponding to the C-terminus of SNAP-25 (NP_003072.2, NP_570824.1).

External Database Links

UniProt:

[P60880](#) [Related reagents](#)

Entrez Gene:

[6616](#) SNAP25 [Related reagents](#)

Synonyms SNAP

RRID AB_808741

Specificity **Goat anti Human SNAP25 antibody** recognizes human synaptosomal-associated protein, 25kD (SNAP-25), also known as Super protein (Sup) or Synaptosomal-associated 25 kDa protein. SNAP-25 is a 206 amino acid, presynaptic plasma membrane protein containing 2 [tSNARE coiled-coil](#) domains and is essential for Ca²⁺-regulated exocytosis of neurotransmitters. SNAP-25 binds to and negatively modulate Ca²⁺ channels, increasing the speed of neurotransmitter release ([Rettig et al. 1996](#)). Goat anti Human SNAP25 antibody detects an epitope within the C-terminal (CT) of human SNAP25 (aa 193-206).

As a result of alternative splicing, SNAP-25 is expressed by neurons and neuroendocrine cells as two predominant isoforms; SNAP-25a and SNAP-25b. SNAP-25a is expressed early in development, whereas SNAP-25b expression is linked to synaptogenesis ([Bark and Wilson 1994](#)). Goat anti human SNAP25 antibody is expected to recognise both isoforms.

SNAP-25 may be a possible therapeutic treatment target for sufferers of hyperactivity disorders ([Barr et al. 2000](#)), whilst SNAP-25 based peptides may be a possible treatment for involuntary muscle spasms ([Hodel 1998](#)).

Western Blotting AHP1124 detects a band of approximately 22kDa in human and mouse brain lysates. A minimum incubation time of 1 hour is recommended with this antibody.

References

- Hodel, A. (1998) SNAP-25. [Int J Biochem Cell Biol. 30 \(10\): 1069-73.](#)
- Lazzarini, R. et al. (2019) Mesenchymal Stem Cells from Nucleus Pulposus and Neural

Differentiation Potential: a Continuous Challenge. [J Mol Neurosci. 67 \(1\): 111-124.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10058 available at: <https://www.bio-rad-antibodies.com/SDS/AHP1124>
10058

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382786:210513'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)