

## Datasheet: AHP1174

<b>Description:</b>	RABBIT ANTI HUMAN INTERLEUKIN-23 (N-TERMINAL)
<b>Specificity:</b>	IL-23 (N-TERMINAL)
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1 - 2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Mouse

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG - liquid

### Antiserum Preparation

Antisera to human IL-23 (NT) were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.

### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.02% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	A 16 amino acid peptide from near the amino terminus of human IL-23.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q9NPF7</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">51561</a>    IL23A    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	SGRF
<b>RRID</b>	AB_844623
<b>Specificity</b>	<p><b>Rabbit anti Human IL-23 antibody</b> recognizes the N-terminal (NT) domain of human interleukin-23 (IL-23), a ~21 kDa heterodimeric cytokine structurally related to IL-12. IL-23 is composed of a p19 subunit encoded by the IL23A gene, and a p40 subunit of interleukin-12 (IL12B). IL-23 is expressed mainly by activated macrophages and dendritic cells, and plays an important role in bridging innate and adaptive responses.</p> <p>IL-23 activates a subset of T cells called Th17 cells, which are characterised by the production of interleukin-17. The IL-23/IL-17 pathway is essential for the induction and maintenance of brain and joint inflammatory autoimmune diseases, including Guillain Barre Syndrome, Multiple sclerosis and rheumatoid arthritis.</p>
<b>Western Blotting</b>	AHP1174 detects a band of approximately 24kDa in mouse pancreas tissue lysate.
<b>Further Reading</b>	<ol style="list-style-type: none"> <li>1. Kikly, K. <i>et al.</i> (2006) The IL-23/Th(17) axis: therapeutic targets for autoimmune inflammation. <a href="#">Curr Opin Immunol. 18 (6): 670-5.</a></li> <li>2. Hu, W. <i>et al.</i> (2006) Interleukin 23 in acute inflammatory demyelination of the peripheral nerve. <a href="#">Arch Neurol. 63 (6): 858-64.</a></li> <li>3. Langrish, C.L. <i>et al.</i> (2004) IL-12 and IL-23: master regulators of innate and adaptive immunity. <a href="#">Immunol Rev. 202: 96-105.</a></li> <li>4. Hunter, C.A. (2005) New IL-12-family members: IL-23 and IL-27, cytokines with divergent functions. <a href="#">Nat Rev Immunol. 5 (7): 521-31.</a></li> </ol>
<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
<b>Guarantee</b>	12 months from date of despatch

---

**Health And Safety Information**      Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/AHP1174>  
10040

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)      [FITC](#)  
Sheep Anti Rabbit IgG (STAR35...)      [RPE](#)  
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)  
Sheep Anti Rabbit IgG (STAR36...)      [DyLight®488](#), [DyLight®680](#), [DyLight®800](#)  
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

### Recommended Useful Reagents

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M382670:210513'

**Printed on 25 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)