

Datasheet: AHP1180 BATCH NUMBER 165216

Description:	RABBIT ANTI IRF7 (C-TERMINAL)
Specificity:	IRF7 (C-TERMINAL)
Other names:	INTERFERON REGULATORY FACTOR 7
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general	protocol r	ecommer	idations, please visit <u>w</u>	ww.bio-		
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry			•			
	Immunohistology - Frozen			•			
	Immunohistology - Paraffin			•			
	ELISA			•			
	Immunoprecipitation			•			
	Western Blotting	•			0.5 - 1.0ug/ml		
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.						
Target Species	Human						
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity a reactivity is derived from personal communications further information.	and workii testing wi	ithin our la	aboratories, peer-revie	wed publications or		
Product Form	Purified IgG - liquid						
Antiserum Prepara	tion Antisera to human IRF7	were raise	ed by rep	eated immunisations o	f rabbits with highly		

purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.02% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	14 amino acid peptide sequence near the carboxy terminus of human IRF7.
External Database Links	UniProt: <u>Q92985</u> <u>Related reagents</u> Entrez Gene: <u>3665</u> IRF7 <u>Related reagents</u>
RRID	AB_844627
Specificity	Rabbit anti IRF7 antibody detects an epitope within the C-terminal (CT) of human interferon regulatory factor 7 (IRF7), a ~54 kDa transcriptional activator of type I interferons (IFNalpha/beta) in response to local and systemic viral infection. IRF7 is mostly expressed by plasmacytoid dendritic cells (pDC), which differentiate in response to IFNs and virus infection. This then stabilizes the IRF7 protein, in turn leading to increased IFN production.
	Multiple IRF7 transcript variants have been identified.
Western Blotting	Multiple IRF7 transcript variants have been identified. AHP1180 detects a band of approximately 51kDa in 293 whole cell lysate.
Western Blotting References	
	 AHP1180 detects a band of approximately 51kDa in 293 whole cell lysate. 1. Rautela, J. <i>et al.</i> (2015) Loss of Host Type-I IFN Signaling Accelerates Metastasis and Impairs NK-cell Antitumor Function in Multiple Models of Breast Cancer. <u>Cancer Immunol Res. 3 (11): 1207-17.</u> 2. Saitoh, S.I. <i>et al.</i> (2017) TLR7 mediated viral recognition results in focal type I interferon
References	 AHP1180 detects a band of approximately 51kDa in 293 whole cell lysate. 1. Rautela, J. <i>et al.</i> (2015) Loss of Host Type-I IFN Signaling Accelerates Metastasis and Impairs NK-cell Antitumor Function in Multiple Models of Breast Cancer. <u>Cancer Immunol Res. 3 (11): 1207-17.</u> 2. Saitoh, S.I. <i>et al.</i> (2017) TLR7 mediated viral recognition results in focal type I interferon secretion by dendritic cells. <u>Nat Commun. 8 (1): 1592.</u> 1. Ning, S. <i>et al.</i> (2005) Regulation of the transcriptional activity of the IRF7 promoter by a pathway independent of interferon signaling. <u>J Biol Chem. 280 (13): 12262-70.</u> 2. Prakash, A. & Levy, D.E. (2006) Regulation of IRF7 through cell type-specific protein
References	 AHP1180 detects a band of approximately 51kDa in 293 whole cell lysate. 1. Rautela, J. <i>et al.</i> (2015) Loss of Host Type-I IFN Signaling Accelerates Metastasis and Impairs NK-cell Antitumor Function in Multiple Models of Breast Cancer. <u>Cancer Immunol Res. 3 (11): 1207-17.</u> 2. Saitoh, S.I. <i>et al.</i> (2017) TLR7 mediated viral recognition results in focal type I interferon secretion by dendritic cells. <u>Nat Commun. 8 (1): 1592.</u> 1. Ning, S. <i>et al.</i> (2005) Regulation of the transcriptional activity of the IRF7 promoter by a pathway independent of interferon signaling. <u>J Biol Chem. 280 (13): 12262-70.</u> 2. Prakash, A. & Levy, D.E. (2006) Regulation of IRF7 through cell type-specific protein stability. <u>Biochem Biophys Res Commun. 342 (1): 50-6.</u> This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1180 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)FITCGoat Anti Rabbit IgG (H/L) (STAR124...)HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRP

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M386148:210519'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint