

## Datasheet: AHP1191

<b>Description:</b>	RABBIT ANTI PAK2 (C-TERMINAL)
<b>Specificity:</b>	PAK2 (C-TERMINAL)
<b>Other names:</b>	p21 ACTIVATED KINASE 2
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			0.5 - 1.0ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Rat, Mouse <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
<b>Product Form</b>	Purified IgG - liquid

**Antiserum Preparation** Antisera to human PAK2 were raised by repeated immunisations of rabbits with highly

purified antigen. Purified IgG prepared by affinity chromatography.

---

<b>Buffer Solution</b>	Phosphate buffered saline
------------------------	---------------------------

---

<b>Preservative Stabilisers</b>	0.02% Sodium Azide
---------------------------------	--------------------

---

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
---------------------------------------	----------------------------

---

<b>Immunogen</b>	14 amino acid peptide sequence near the carboxy terminus of human PAK2.
------------------	---

---

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q13177</a> <a href="#">Related reagents</a>
	<b>Entrez Gene:</b> <a href="#">5062</a> PAK2 <a href="#">Related reagents</a>

---

<b>RRID</b>	AB_844634
-------------	-----------

---

<b>Specificity</b>	<p><b>Rabbit anti Human PAK2 antibody</b> recognizes p21-activated kinase-2 (PAK2), a ubiquitously expressed, ~64 kDa, serine/threonine protease found at higher levels in skeletal muscle, ovary, thymus and spleen. p21-activated kinases (PAKs) bind activated Cdc42 and Rac via a Cdc42/Rac interactive binding (CRIB) domain. Group 1 PAK kinases (PAK1, 2 and 3) regulate the activity of an adjacent conserved autoinhibitory domain and are associated with cellular processes, including cytoskeleton rearrangement, apoptosis and MAP kinase signalling. Group 2 PAKs (PAK4, 5, and 6) lack the autoinhibitory domain and are not activated by Cdc42/Rac binding.</p> <p>PAK2 has a cytostatic effect on cells and is activated in response to cellular stresses such as hyperosmolarity, serum starvation, contact inhibition and DNA damage from radiation and chemicals. Unlike other PAKs, PAK2 can be activated by proteolytic cleavage to generate a constitutively active fragment, PAK2p34, which induces a cell death response. This induction is thought to be regulated by Rho GTPase-activating protein 10 which inhibits the protein kinase activity of PAK2p34 and causes it to localise to the perinuclear region instead of the nucleus. PAK2 may also play a role in X-linked mental retardation.</p>
--------------------	--

---

<b>Western Blotting</b>	AHP1191 detects a band of approximately 58kDa in Jurkat cell lysates.
-------------------------	---

---

<b>Further Reading</b>	<ol style="list-style-type: none"><li>1. Roig, J. &amp; Traugh, J.A. (2001) Cytostatic p21 G protein-activated protein kinase gamma-PAK. <a href="#">Vitam Horm. 62: 167-98.</a></li><li>2. Rudel, T. &amp; Bokoch, G.M. (1997) Membrane and morphological changes in apoptotic cells regulated by caspase-mediated activation of PAK2. <a href="#">Science. 276 (5318): 1571-4.</a></li><li>3. Jaffer, Z.M. &amp; Chernoff, J. (2002) p21-activated kinases: three more join the Pak. <a href="#">Int J Biochem Cell Biol. 34 (7): 713-7.</a></li></ol>
------------------------	--

---

<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for
----------------	---

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)  
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)  
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)  
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)  
Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@680](#), [DyLight@800](#)

### Recommended Useful Reagents

- [ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)  
[100x ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)  
[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382793:210513'

Printed on 18 May 2021