Datasheet: AHP1233 BATCH NUMBER 166840

Description:	RABBIT ANTI GSK3 BETA (pSer9)
Specificity:	GSK3 BETA (pSer9)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry			-				
	Immunohistology - Frozen			•				
	Immunohistology - Paraffin			•				
	ELISA			•				
	Western Blotting	-			1/1000			
	Where this product has n	ot been te	ested for u	se in a particular tech	inique this does not			
	necessarily exclude its us	se in such	procedur	es. Suggested workin	g dilutions are given as			
	a guide only. It is recomn system using appropriate			•	or use in their own			
Target Species	Rat							
Species Cross Reactivity	 Based on sequence similarity, is expected to react with:Mouse, Dog, Primate, Xenopus, Human, Bovine, Chicken, Zebrafish N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. 							
Product Form	Purified IgG - liquid							
Antiserum Preparatio	n Antisera to rat GSK3 beta highly purified antigen. P	,		• •				
Buffer Solution	10mM Hepes pH7.5							

Stabilisers 5000 Second Stream Packet (Naty) Stabilisers 50% Glycerol 0.01% Bovine Serum Albumin Immunogen Phosphopeptide corresponding to amino acid sequence surrounding phoshorylated seine 9 of rat GSK3 beta. External Database Links P18266 RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is sexpressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mittogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. <u>Biochem J. 303:701-4</u> . 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulat		
b3% Gycerol 0.01% Bovine Serum Albumin Immunogen Phosphopeptide corresponding to amino acid sequence surrounding phoshorylated seine 9 of rat GSK3 beta. External Database Links UniProt: P18266 Related reagents Entrez Gene: 84027 Gsk3b teta Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-proteir kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the tests, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. <i>et al.</i> (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and st	Preservative	0.09% Sodium Azide (NaN ₃)
Immunogen Phosphopeptide corresponding to amino acid sequence surrounding phoshorylated seine 9 of rat GSK3 beta. External Database UniProt: P18266 Related reagents Entrez Gene: 84027 Supcificity Rabbit anti Rat GSK3 beta (pSer0) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting A.HP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. <i>et al.</i> (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8, 3. Green, J.B. (2004) LkD1 and GSK3 beta: are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store al ~20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-	Stabilisers	50% Glycerol
9 of rat GSK3 beta. External Database Links UniProt: P18266 Related reagents Entrez Gene: 8402Z GsK3b RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Blochem, J. 303:701-4. 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Blochem, Blophys, Res. Commun, 247:851-8. 3. Green, J.B. (2004) LkD1 and GSK3-beta: kinases at t		0.01% Bovine Serum Albumin
Links UniProt: P18266 Related reagents Entrez Gene: 84027 Skip Related reagents RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. et al. (1994) WhT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem, Biophys. Res. Commun. 247:851-8. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliq	Immunogen	
P18266 Related reagents Entrez Gene: 84027 GsX3b Related reagents RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. <i>et al.</i> (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem. J. 303:701-4. 2. Papkoff, J. <i>et al.</i> (1994) WhT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-3. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycle. 3 (1): 12-4. 4. Garrido, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon	External Database	
Entrez Gene: B4027 Gsk3b Related reagents RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. <i>et al.</i> (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. <i>et al.</i> (1994) WIT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycle. 3 (1): 12-4. 4. Garrido, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storag	Links	
84027 Gsk3b Related reagents RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem. J. 303:701-4. 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycie. 3 (1): 12-4. 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product i		P18266 Related reagents
RRID AB_877475 Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycice. 3 (1): 12-4. 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store al -20°C on receipt		Entrez Gene:
Specificity Rabbit anti Rat GSK3 beta (pSer9) antibody recognizes GSK3 beta, also known as glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. 1. Stambolic, V. <i>et al.</i> (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. <u>Biochem, J. 303:701-4</u> . 2. Papkoff, J. <i>et al.</i> (1994) WIT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. <u>Biochem, Biophys. Res. Commun. 247:851-8</u> . 3. Green, J. B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. <u>Cell Cycle. 3 (1): 12-4</u> . 4. Garrido, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. <u>FEBS Lett. 581:1579-86</u> . Storage This product is shipped at ambient temperature. It is recommended to aliquot and store al -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		84027 Gsk3b Related reagents
glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at low levels in lung, brain and kidney tissue. GSK3 beta is involved in many signalling pathways including energy metabolism, body pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. <u>Biochem. J. 303:701-4.</u> 2. Papkoff, J. et al. (1994) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. <u>Biochem. J. 303:701-4.</u> 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycle. 3 (1): 12-4. 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store al -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.	RRID	AB_877475
pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several regulatory proteins including glycogen synthase. Phosphorylation of serine-9 inhibits GSK3 beta function. Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycle. 3 (1): 12-4. 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in	Specificity	glycogen synthase kinase-3 beta (GSK-3 beta), factor A (FA) and serine/threonine-protein kinase GSK3B, when phosphorylated at serine 9. GSK3 is a ~46 kDa member of the serine/threonine protein kinase family. Two isforms of GSK3 exist, alpha and beta. GSK3 beta is expressed in the testis, thymus, prostate, and ovary as well as being expressed at
Western Blotting AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates. Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. Biochem, J. 303:701-4. 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. Biochem. Biophys. Res. Commun. 247:851-8. 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. Cell Cycle. 3 (1): 12-4. 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. FEBS Lett. 581:1579-86. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in		pattern formation and microtubule growth and stability (which aids axon growth). GSK3 beta is also thought to be involved in oncogenesis and the hormonal control of several
Further Reading 1. Stambolic, V. et al. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. <u>Biochem, J. 303:701-4.</u> 2. Papkoff, J. et al. (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. <u>Biochem. Biophys. Res. Commun. 247:851-8.</u> 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. <u>Cell Cycle. 3 (1): 12-4.</u> 4. Garrido, J. et al. (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. <u>FEBS Lett. 581:1579-86.</u> Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in		Phosphorylation of serine-9 inhibits GSK3 beta function.
 intact cells via serine 9 phosphorylation. <u>Biochem, J. 303:701-4.</u> 2. Papkoff, J. <i>et al.</i> (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. <u>Biochem. Biophys. Res. Commun. 247:851-8.</u> 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. <u>Cell Cycle. 3 (1): 12-4.</u> 4. Garrido, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon formation. <u>FEBS Lett. 581:1579-86.</u> Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. 	Western Blotting	AHP1233 detects a band of approximately 46kDa in rat cortex cell lysates.
-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in	Further Reading	 intact cells via serine 9 phosphorylation. <u>Biochem, J. 303:701-4.</u> 2. Papkoff, J. <i>et al.</i> (1998) WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. <u>Biochem. Biophys. Res. Commun. 247:851-8.</u> 3. Green, J.B. (2004) Lkb1 and GSK3-beta: kinases at the center and poles of the action. <u>Cell Cycle. 3 (1): 12-4.</u> 4. Garrido, J. <i>et al.</i> (2007) GSK3 alpha and GSK3 beta are necessary for axon formation.
short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in	Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at

Sheep Anti Rabbit IgG (STAR34...)FITCGoat Anti Rabbit IgG (H/L) (STAR124...)HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio	-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M390260:210825'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint