

## Datasheet: AHP1274

<b>Description:</b>	RABBIT ANTI HUMAN DELTA-LIKE PROTEIN 4
<b>Specificity:</b>	DELTA-LIKE PROTEIN 4
<b>Other names:</b>	DLL4
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin	▪			5 - 15ug/ml
ELISA	▪			1/3000 - 1/15000
Western Blotting	▪			1/300 - 1/1500
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Based on sequence similarity, is expected to react with: Mouse, Rat, Chimpanzee  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG - liquid

### Antiserum Preparation

Antisera to DLL4 were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG was prepared by affinity chromatography.

### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml
<b>Immunogen</b>	Synthetic peptide corresponding to the internal region of human DLL4.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q9NR61</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">54567</a>    DLL4    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_2092966
<b>Specificity</b>	<p><b>Rabbit anti Human Delta-like Protein 4 antibody</b> detects human Delta-like protein 4 (DLL4), also known as Delta-4. DLL4 is a 685 amino acid single pass type 1 transmembrane glycoprotein of ~72 kDa containing a single <a href="#">DSL</a> domain and eight <a href="#">EGF-like</a> domains. Human DLL4 is the homologue of the Drosophila delta protein, and functions as a transmembrane bound ligand to the Notch receptor, Notch1.</p> <p>DLL4 is expressed in the vasculature and plays a critical role in vascular development. It is induced by vascular endothelial growth factor (VEGF), as a negative feedback regulator to regulate angiogenic sprouting and promote the formation of a differentiated vascular network (<a href="#">Mailhos et al. 2001</a>). DLL4 is strongly expressed in tumor vessels of primary renal tumors (<a href="#">Patel et al. 2005</a>), bladder cancer (<a href="#">Patel et al. 2006</a>), and inhibition of DLL4 results in increased vascular proliferation but defective maturation. This in turn leads to a decrease in tumor growth, with no apparent toxicity (<a href="#">Ridgway et al. 2006</a>).</p> <p>Expression of DLL4 in normal human dermis is low in foetal tissues, becomes more intense during early life (0-20 years) and gradually declines thereafter as shown by immunohistochemistry on FFPE tissues using rabbit anti human DLL4 antibody (<a href="#">Gunin et al. 2014</a>).</p>
<b>Histology Positive Control Tissue</b>	Human ovary.
<b>Western Blotting</b>	AHP1274 detects a band of approximately 74kDa in pancreatic cell lysates.
<b>References</b>	<ol style="list-style-type: none"> <li>Villaamil, V.M. <i>et al.</i> (2012) Multiple biomarker tissue arrays: A computational approach to identifying protein-protein interactions in the EGFR/ERK signalling pathway. <a href="#">J Mol Signal. 7: 14.</a></li> <li>Medina Villaamil, V. <i>et al.</i> (2012) Searching for Hif1-<math>\alpha</math> interacting proteins in renal cell carcinoma. <a href="#">Clin Transl Oncol. 14: 698-708.</a></li> <li>You, C. <i>et al.</i> (2013) Loss of CCM3 impairs DLL4-Notch signalling: implication in endothelial angiogenesis and in inherited cerebral cavernous malformations. <a href="#">J Cell Mol Med. 17 (3): 407-18.</a></li> </ol>

4. El Hindy, N. *et al.* (2013) Implications of Dll4-Notch signaling activation in primary glioblastoma multiforme. [Neuro Oncol. 15: 1366-78.](#)
5. Gunin, A.G. *et al.* (2014) Age-related changes in angiogenesis in human dermis. [Exp Gerontol. 55C: 143-51.](#)
6. Hjelmgren O *et al.* (2016) Increased Vascularization in the Vulnerable Upstream Regions of Both Early and Advanced Human Carotid Atherosclerosis. [PLoS One. 11 \(12\): e0166918.](#)

---

**Further Reading** 1. Thurston, G. *et al.* (2007) The Delta paradox: DLL4 blockade leads to more tumour vessels but less tumour growth. [Nat Rev Cancer. 7 \(5\): 327-31.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/AHP1274>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)  
 Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)  
 Sheep Anti Rabbit IgG (STAR35...) [RPE](#)  
 Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)  
[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
 'M389130:210806'

Printed on 19 Jan 2024