Datasheet: AHP1278 BATCH NUMBER 151215

Description:	RABBIT ANTI HUMAN CTGF
Specificity:	CTGF
Other names:	CONNECTIVE TISSUE GROWTH FACTOR
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is
	derived from testing within our laboratories, peer-reviewed publications or personal
	communications from the originators. Please refer to references indicated for further
	information. For general protocol recommendations, please visit <u>www.bio-</u>
	rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen				
Immunohistology - Paraffin (1)	•			
ELISA	-			0.5 - 2.0ug/ml
Western Blotting	-			0.1 - 0.2ug/ml
Immunofluorescence				
Functional Assays	-			6.0 - 9.0ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Product Form	Purified IgG - lyophilised
Reconstitution	Reconstitute with 0.1ml distilled water. For long term storage the addition of 0.09% sodium azide is recommended. NB: For functional studies do not add azide.

Antiserum Preparation Antisera to human CTGF were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Dheenhote huffered coline				
Buller Solution	Phosphate buffered saline				
Preservative Stabilisers	None present				
Carrier Free	Yes				
Approx. Protein Concentrations	IgG concentration 1.0mg/ml after reconstitution.				
Immunogen	Recombinant human CTGF.				
External Database Links	UniProt: <u>P29279</u> <u>Related reagents</u> Entrez Gene: <u>1490</u> CTGF <u>Related reagents</u>				
Synonyms	CCN2, HCS24, IGFBP8				
RRID	AB_2087370				
Specificity	 Rabbit anti Human CTGF antibody recognizes human CTGF (connective tissue growth factor), a 323 amino acid ~38 kDa polypeptide member of the CCN family of secreted cysteine rich regulatory proteins. CTGF is produced by vascular endothelial and umbilical vein cells, and interacts with several other growth factors, including transforming growth factor beta (TGF-beta), vascular endothelial growth factor (VEGF) and bone morphogenetic proteins (BMPs). CTGF is a heparin-binding pleiotropic growth factor involved in chondrogenesis and potentially skeletogenesis. CTGF promotes the adhesion of epithelial cells and fibroblasts, stimulates integrin expression, and also promotes endothelial cell survival, migration and adhesion, implicating CTGF in angiogenesis. CTGF induces extracellular matrix formation, including the deposition of excessive collagen in fibrotic diseases, and has also been implicated in wound healing. 				
ELISA	This purified human CTGF antibody may ne used in an indirect ELISA or as the capture reagent in a sandwich ELISA with our <u>biotinylated human CTGF antibody</u> (AHP1278B) as the detection reagent and <u>recombinant human CTGF</u> (PHP174) as the standard.				
References	 Waddell, J.M. <i>et al.</i> (2011) CTGF expression is up-regulated pregnancy and influences HTR-8/Svneo cell adhesion and netw <u>Reprod. 26 (1): 67-75.</u> Westermann, D. <i>et al.</i> (2009) Gene deletion of the kinin receiption inflammation and fibrosis during the development of experiment cardiomyopathy. <u>Diabetes. 58: 1373-81.</u> Vasilieva, O.V. <i>et al.</i> (2016) Connective tissue growth factor dermis through ontogenesis. <u>Russian Journal of Developmenta</u> 	work formation. <u>Hum</u> eptor B1 attenuates cardiac ntal diabetic (CTGF) in the human			

Further Reading	1. Moussad, E.E. & Brigstock, D.R. (2000) Connective tissue growth factor: what's in a name? <u>Mol Genet Metab. 71 (1-2): 276-92.</u>
Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C.
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/AHP1278 10162
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34)	<u>FITC</u>			
Goat Anti Rabbit IgG (Fc) (STAR121)	Biotin, FITC, HRP			
Sheep Anti Rabbit IgG (STAR35)	RPE			
Goat Anti Rabbit IgG (H/L) (STAR124) <u>HRP</u>				

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376 Worldv	wide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372478:200706'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint