

# Datasheet: AHP1278 BATCH NUMBER 155862

Description:	RABBIT ANTI HUMAN CTGF
Specificity:	CTGF
Other names:	CONNECTIVE TISSUE GROWTH FACTOR
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)	•			
ELISA	•			0.5 - 2.0ug/ml
Western Blotting	•			0.1 - 0.2ug/ml
Immunofluorescence	•			
Functional Assays	•			6.0 - 9.0ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - lyophilised	
Reconstitution	Reconstitute with 0.1ml distilled water. For long term storage the azide is recommended. NB: For functional studies do not add	

**Antiserum Preparation** Antisera to human CTGF were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml after reconstitution.
Immunogen	Recombinant human CTGF.
External Database Links	UniProt: P29279 Related reagents  Entrez Gene: 1490 CTGF Related reagents
Synonyms	CCN2, HCS24, IGFBP8
RRID	AB_2087370
Specificity	Rabbit anti Human CTGF antibody recognizes human CTGF (connective tissue growth factor), a 323 amino acid ~38 kDa polypeptide member of the CCN family of secreted cysteine rich regulatory proteins. CTGF is produced by vascular endothelial and umbilical vein cells, and interacts with several other growth factors, including transforming growth factor beta (TGF-beta), vascular endothelial growth factor (VEGF) and bone morphogenetic proteins (BMPs).  CTGF is a heparin-binding pleiotropic growth factor involved in chondrogenesis and potentially skeletogenesis. CTGF promotes the adhesion of epithelial cells and fibroblasts, stimulates integrin expression, and also promotes endothelial cell survival, migration and adhesion, implicating CTGF in angiogenesis. CTGF induces extracellular matrix formation, including the deposition of excessive collagen in fibrotic diseases, and has also been implicated in wound healing.
ELISA	This purified human CTGF antibody may ne used in an indirect ELISA or as the capture reagent in a sandwich ELISA with our <u>biotinylated human CTGF antibody</u> (AHP1278B) as the detection reagent and <u>recombinant human CTGF</u> (PHP174) as the standard.
References	<ol> <li>Waddell, J.M. <i>et al.</i> (2011) CTGF expression is up-regulated by PROK1 in early pregnancy and influences HTR-8/Svneo cell adhesion and network formation. <u>Hum Reprod. 26 (1): 67-75.</u></li> <li>Westermann, D. <i>et al.</i> (2009) Gene deletion of the kinin receptor B1 attenuates cardiac inflammation and fibrosis during the development of experimental diabetic cardiomyopathy. <u>Diabetes. 58: 1373-81.</u></li> <li>Vasilieva, O.V. <i>et al.</i> (2016) Connective tissue growth factor (CTGF) in the human dermis through ontogenesis. <u>Russian Journal of Developmental Biology. 47 (2): 63-68.</u></li> </ol>

Further Reading	1. Moussad, E.E. & Brigstock, D.R. (2000) Connective tissue growth factor: what's in name? Mol Genet Metab. 71 (1-2): 276-92.		
Storage	Prior to reconstitution store at -20°C.		
	After reconstitution store at -20°C.		
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.		
Guarantee	12 months from date of despatch		
Health And Safety	Material Safety Datasheet documentation #10162 available at:		
Information	https://www.bio-rad-antibodies.com/SDS/AHP1278 10162		
Regulatory	For research purposes only		

## Related Products

## **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) FITC

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

## **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)
TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372478:200706'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint