Datasheet: AHP1294 BATCH NUMBER 160332

Description:	RABBIT ANTI YEAST RAD9
Specificity:	RAD9
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry							
	Immunohistology - Frozen			-				
	Immunohistology - Paraffin			-				
	ELISA	-			1/2000 - 1/4000			
	Immunoprecipitation			•				
	Western Blotting	•			1/50 - 1/250			
	Functional Assays			•				
	Where this product has r	not been te	ested for u	se in a particular teo	chnique this does not			
	necessarily exclude its us	se in such	procedure	es. Suggested worki	ing dilutions are given as			
	a guide only. It is recomn system using appropriate			•	t for use in their own			
Target Species	Yeast							
Product Form	Purified IgG - liquid							
Antiserum Preparatio	n Antisera to yeast Rad9 w purified antigen. Purified				• •			
Buffer Solution	Phosphate buffered salin	е						
Preservative Stabilisers	<0.1% Sodium Azide (Na	aN ₃)						
Approx. Protein	IgG concentration 0.5mg	/ml						

Concentrations

DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA rep Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase genomic instability.Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated a non-phosphorylated forms of this protein.Western BlottingAHP1294 detects a band of approximately 60kDa.Further Reading1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 prot in sensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6. 2. Sweeney, F.D. et al. (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adaptor allow Rad53 activation. Curr Biol. 15 (15): 1364-75.StorageThis product is shipped at ambient temperature. It is recommended to aliquot and store	Immunogen	Synthetic peptide corresponding to aa 1125-1139 of yeast Rad9 protein conjugated to Keyhole Limpet Haemocyanin (KLH).				
P14737 Related reagents RRID AB_906114 Specificity Rabbit anti yeast Rad9 antibody detects Saccharomyces cerevisiae Rad9, a protein involved in the DNA damage checkpoint. In <i>S. cerevisiae</i> this checkpoint is mainly controlled by the PI(3) kinase-like kinase (PIKK) Mec1, which is activated in response: DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA rep Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase genomic instability. Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated a non-phosphorylated forms of this protein. Western Blotting AHP1294 detects a band of approximately 60kDa. Further Reading 1. Toh, G. W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 prot in sensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6. 2. Sweeney, F.D. <i>et al.</i> (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapte allow Rad53 activation. Curr Biol. 15 (15): 1364-75. Storage This product is shipped at ambient temperature. It is recommended to aliquot and storn -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Augental Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040 </th <th>External Database</th> <th></th>	External Database					
RRID AB_906114 Specificity Rabbit anti yeast Rad9 antibody detects Saccharomyces cerevisiae Rad9, a protein involved in the DNA damage checkpoint. In S. cerevisiae this checkpoint is mainly controlled by the PI(3) kinase-like kinase (PIKK) Mec1, which is activated in response DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA rep. Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase genomic instability. Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated a non-phosphorylated forms of this protein. Western Blotting AHP1294 detects a band of approximately 60kDa. Further Reading 1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 profin in sensing and responding to DNA damage. Biochem Soc Trans. 31 (P1 1): 242-6, 2. Sweeney, F.D. et al. (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapte allow Rad53 activation. Curr Biol. 15 (15): 1364-75. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: https	Links	UniProt:				
Specificity Rabbit anti yeast Rad9 antibody detects Saccharomyces cerevisiae Rad9, a protein involved in the DNA damage checkpoint. In S. cerevisiae this checkpoint is mainly controlled by the PI(3) kinase-like kinase (PIKK) Mec1, which is activated in response DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA rep Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase genomic instability. Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated a non-phosphorylated forms of this protein. Western Blotting AHP1294 detects a band of approximately 60kDa. Further Reading 1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 prot in sensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6, 2. Sweeney, F.D. et al. (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapter allow Rad53 activation. Curr Biol. 15 (15): 1364-75. Storage This product is shipped at ambient temperature. It is recommended to aliquot and storr -20°C or receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: htttps://www.bio-rad-antibodies.com/SDS/AHP1294<		P14737 Related reagents				
involved in the DNA damage checkpoint. In S. cerevisiae this checkpoint is mainly controlled by the PI(3) kinase-like kinase (PIKK) Mec1, which is activated in response DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA rep Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase genomic instability. Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated a non-phosphorylated forms of this protein. Western Blotting AHP1294 detects a band of approximately 60kDa. Further Reading 1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 prot in sensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6, 2. Sweeney, F.D. <i>et al.</i> (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapte allow Rad53 activation. Curr Biol. 15 (15): 1364-75. Storage This product is shipped at ambient temperature. It is recommended to aliquot and stor -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040	RRID	AB_906114				
Non-phosphorylated forms of this protein.Western BlottingAHP1294 detects a band of approximately 60kDa.Further Reading1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 prot in sensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6. 2. Sweeney, F.D. <i>et al.</i> (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapted allow Rad53 activation. Curr Biol. 15 (15): 1364-75.StorageThis product is shipped at ambient temperature. It is recommended to aliquot and store -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040	Specificity	involved in the DNA damage checkpoint. In <i>S. cerevisiae</i> this checkpoint is mainly controlled by the PI(3) kinase-like kinase (PIKK) Mec1, which is activated in response to DNA damage. Activated Mec1 hyperphosphorylates Rad9, which in turn triggers the binding of Rad53, the orthologue of mammalian Chk2. Activated Rad53 specifically targets substrates required for cell cycle arrest, gene expression and efficient DNA repair. Mutations affecting Rad9 impair checkpoint induced cell-cycle arrest and increase				
Further Reading 1. Toh, G.W. & Lowndes, N.F. (2003) Role of the Saccharomyces cerevisiae Rad9 profinsensing and responding to DNA damage. Biochem Soc Trans. 31 (Pt 1): 242-6. 2. Sweeney, F.D. et al. (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapter allow Rad53 activation. Curr Biol. 15 (15): 1364-75. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040		Rabbit anti yeast Rad9 antibody (AHP1294) is reactive with both the phosphorylated and non-phosphorylated forms of this protein.				
in sensing and responding to DNA damage. <u>Biochem Soc Trans. 31 (Pt 1): 242-6.</u> 2. Sweeney, F.D. <i>et al.</i> (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adapte allow Rad53 activation. <u>Curr Biol. 15 (15): 1364-75.</u> StorageThis product is shipped at ambient temperature. It is recommended to aliquot and store -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040	Western Blotting	AHP1294 detects a band of approximately 60kDa.				
-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040	Further Reading	2. Sweeney, F.D. et al. (2005) Saccharomyces cerevisiae Rad9 acts as a Mec1 adaptor to				
frost-free freezers is not recommended. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1294 10040	Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
Health And Safety Material Safety Datasheet documentation #10040 available at: Information <u>https://www.bio-rad-antibodies.com/SDS/AHP1294</u> 10040						
Information <u>https://www.bio-rad-antibodies.com/SDS/AHP1294</u> 10040	Guarantee	12 months from date of despatch				
Regulatory For research purposes only		https://www.bio-rad-antibodies.com/SDS/AHP1294				
	Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

 Sheep Anti Rabbit IgG (STAR34...)
 FITC

 Sheep Anti Rabbit IgG (STAR35...)
 RPE

 Goat Anti Rabbit IgG (Fc) (STAR121...)
 Biotin, FITC, HRP

 Sheep Anti Rabbit IgG (STAR36...)
 DyLight®488, DyLight®680, DyLight®800

 Goat Anti Rabbit IgG (H/L) (STAR124...)
 HRP

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389132:210806'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint