

# Datasheet: AHP1391 BATCH NUMBER 148911

| Description:     | GOAT ANTI HUMAN Ku80 |  |
|------------------|----------------------|--|
| Specificity:     | Ku80                 |  |
| Other names:     | XRCC5                |  |
| Format:          | Purified             |  |
| Product Type:    | Polyclonal Antibody  |  |
| Isotype:         | Polyclonal IgG       |  |
| Quantity: 0.1 mg |                      |  |

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry             |     |    |                |                    |
| Immunohistology - Frozen   |     |    |                |                    |
| Immunohistology - Paraffin |     |    |                |                    |
| ELISA                      | •   |    |                | 1/16,000           |
| Immunoprecipitation        |     |    |                |                    |
| Western Blotting           | •   |    |                | 0.1 - 0.3ug/ml     |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human                 |
|----------------|-----------------------|
| Product Form   | Purified IgG - liquid |

**Antiserum Preparation** Antiserum to human Ku80 was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

| Buffer Solution             | TRIS buffered saline  |
|-----------------------------|---|
| Preservative<br>Stabilisers | 0.02% Sodium Azide (NaN <sub>3</sub> )<br>0.5% Bovine Serum Albumin |
| Approx. Protein             | IgG concentration 0.5mg/ml  |

| Concentrations                   |   |
|----------------------------------|---|
| Immunogen                        | Peptide sequence C-SLAKKDEKTDTLED from the internal region of Ku80 (NP_066964.1).   |
| External Database<br>Links       | UniProt:  P13010 Related reagents  Entrez Gene:  7520 XRCC5 Related reagents  |
| Synonyms                         | G22P2   |
| RRID                             | AB_2218624  |
| Specificity                      | <b>Goat anti Human Ku80 antibody</b> recognizes the 80 kDa subunit of human Ku protein, an evolutionarily conserved nuclear ATP-dependent DNA helicase, involved in a major proportion of DNA repair and in V(D)J recombination.  |
|                                  | The Ku protein, originally described as an autoantigen, exists as a tightly associated heterodimer consisting of a 70 kDa (Ku70) and 80 kDa (Ku80) subunit which binds to DNA double-strand break ends (DSB). DNA bound Ku recruits the large catalytic subunit DNA-PKcs to form the DNA-dependent protein kinase complex DNA-PK, facilitating DNA repair by the non-homologous end-joining (NHEJ) pathway.   |
| Western Blotting                 | AHP1391 detects a band of approximately 80-85kDa in HeLa cell lysates. An additional 75kDa band of unknown origin may be observed.  |
| References                       | <ol> <li>Koike, M. (2002) Dimerization, translocation and localization of Ku70 and Ku80 proteins. <u>J Radiat Res. 43 (3): 223-36.</u></li> <li>Dedieu, S. <i>et al.</i> (2010) The cytoprotective drug amifostine modifies both expression and activity of the pro-angiogenic factor VEGF-A. <u>BMC Med. 8:19.</u></li> <li>Slupianek, A. <i>et al.</i> (2011) BCR/ABL Stimulates WRN to Promote Survival and Genomic Instability. <u>Cancer Res. 71: 842-51.</u></li> </ol> |
| Storage                          | Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.   |
| Guarantee                        | 12 months from date of despatch   |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10058 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP1391">https://www.bio-rad-antibodies.com/SDS/AHP1391</a> 10058  |
| Regulatory                       | For research purposes only  |

# **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363917:200529'

### Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint