

Datasheet: AHP1440 BATCH NUMBER 155950

Description:	RABBIT ANTI HUMAN PTK7 (N-TERMINAL)
Specificity:	PTK7 (N-TERMINAL)
Other names:	CCK-4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been re	ported to wo	ork in th	e following applicatio	ons. This information is	
	derived from testing withi	n our labora	tories, I	peer-reviewed publication	ations or personal	
	communications from the originators. Please refer to references indicated for further					
		information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/proto		onninon			
	Tad-antibodies.com/proto		No	Not Determined	Suggested Dilution	
	Flow Cytometry	162	NO	Not Determined	Suggested Dilution	
	Immunohistology - Frozen			-		
				-		
	Immunohistology - Paraffin (1)	•				
	ELISA			-		
	Immunoprecipitation			-		
	Western Blotting	-		-	0.5 - 1.0ug/ml	
	-	-				
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its us a guide only. It is recomm	nended that	rocedur the use	es. Suggested working the antibody	ng dilutions are given as	
	•	nended that negative/pc s antigen re	rocedur the use ositive c etrieval	es. Suggested working of titrates the antibody ontrols. Using heat treatme	ng dilutions are given as y for use in their own nt prior to staining of	
Target Species	a guide only. It is recomm system using appropriate (1) This product require	nended that negative/pc s antigen re	rocedur the use ositive c etrieval	es. Suggested working of titrates the antibody ontrols. Using heat treatme	ng dilutions are given as y for use in their own nt prior to staining of	
	a guide only. It is recomm system using appropriate (1) This product require paraffin sections. Sodiu	nended that negative/pc s antigen re	rocedur the use ositive c etrieval	es. Suggested working of titrates the antibody ontrols. Using heat treatme	ng dilutions are given a y for use in their own nt prior to staining of	
Species Cross	a guide only. It is recomm system using appropriate (1) This product requires paraffin sections. Sodiu Human Reacts with: Rat, Mouse	nended that negative/po s antigen re im citrate b	rocedur the use ositive c etrieval	es. Suggested working ontrols. Using heat treatme H 6.0 is recommend	ng dilutions are given a y for use in their own nt prior to staining of led for this purpose.	
Species Cross	a guide only. It is recomm system using appropriate (1) This product requires paraffin sections. Sodiu Human Reacts with: Rat, Mouse N.B. Antibody reactivity a	nended that negative/pc s antigen re um citrate b	rocedur the use ositive c etrieval ouffer p	es. Suggested workin or titrates the antibody controls. Using heat treatme H 6.0 is recommend ons may vary betwee	ng dilutions are given a y for use in their own nt prior to staining of led for this purpose. n species. Cross	
Species Cross	a guide only. It is recomm system using appropriate (1) This product requires paraffin sections. Sodiu Human Reacts with: Rat, Mouse N.B. Antibody reactivity a reactivity is derived from	nended that negative/po s antigen re im citrate b and working testing within	rocedur the use ositive c etrieval ouffer p condition	es. Suggested working on titrates the antibody ontrols. Using heat treatme H 6.0 is recommend ons may vary betwee boratories, peer-revie	ng dilutions are given a y for use in their own nt prior to staining of led for this purpose. n species. Cross ewed publications or	
Target Species Species Cross Reactivity	a guide only. It is recomm system using appropriate (1) This product requires paraffin sections. Sodiu Human Reacts with: Rat, Mouse N.B. Antibody reactivity a reactivity is derived from personal communications	nended that negative/po s antigen re im citrate b and working testing within	rocedur the use ositive c etrieval ouffer p condition	es. Suggested working on titrates the antibody ontrols. Using heat treatme H 6.0 is recommend ons may vary betwee boratories, peer-revie	ng dilutions are given a y for use in their own nt prior to staining of led for this purpose. n species. Cross ewed publications or	
Species Cross	a guide only. It is recomm system using appropriate (1) This product requires paraffin sections. Sodiu Human Reacts with: Rat, Mouse N.B. Antibody reactivity a reactivity is derived from	nended that negative/po s antigen re im citrate b and working testing within	rocedur the use ositive c etrieval ouffer p condition	es. Suggested working on titrates the antibody ontrols. Using heat treatme H 6.0 is recommend ons may vary betwee boratories, peer-revie	ng dilutions are given a y for use in their own nt prior to staining of led for this purpose. n species. Cross ewed publications or	

Preparation	Antisera to human PTK7 (NT) were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared by affinity chromatography.				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.02% Sodium Azide				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				
Immunogen	A 15 amino acid peptide located near the N-terminal region of human PTK7.				
External Database Links	UniProt: Q13308 Related reagents Entrez Gene: 5754 PTK7 Related reagents				
Synonyms	CCK4				
RRID	AB_2174361				
Specificity	Rabbit anti Human PTK7 antibody recognizes the N-terminal region of human Protein tyrosine kinase-like 7 (PTK7), a member of the receptor tyrosine kinase family, otherwise known as Colon Carcinoma Kinase 4 (CCK4). PTK7 is highly expressed in the kidney, liver, lung, pancreas and melanocytes with weaker expression being detected in the thyroid gland, ovaries, brain, heart and skeletal muscle.				
	PTK7 may play a role in embryonic development. It is thought to function as a cell adhesion molecule and as a novel regulator of the planar cell polarity (PCP) pathway in vertebrates.				
	PTK7 may be involved in the tumour progression. A loss of PTK7 expression has been observed in melanoma cell lines, whereas increased levels of PTK7 have been detected in high-throughput acute myeloid leukaemia samples. In addition, PTK7 is not expressed in normal colon but has been detected in colon carcinomas and as such is a potential marker for tumour progression.				
Western Blotting	AHP1440 detects a band of approximately 86kDa in colon tissue lysates.				
Further Reading	 Lu, X. <i>et al.</i> (2004) PTK7/CCK-4 is a novel regulator of planar cell polarity in vertebrates. <u>Nature. 430 (6995): 93-8.</u> Easty, D.J. <i>et al.</i> (1997) Loss of expression of receptor tyrosine kinase family genes PTK7 and SEK in metastatic melanoma. <u>Int J Cancer. 71 (6): 1061-5.</u> 				
Storage	Store at +4°C or at -20°C if preferred.				

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1440 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)FITCGoat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (H/L) (STAR124...)HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363930:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint