

## Datasheet: AHP1484

<b>Description:</b>	RABBIT ANTI BRCC36
<b>Specificity:</b>	BRCC36
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			0.5 - 1.0ug/ml
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
<b>Product Form</b>	Purified IgG - liquid

**Antiserum Preparation** Antiserum to human BRCC36 was raised by repeated immunisation of rabbits with highly

purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

---

<b>Buffer Solution</b>	Phosphate buffered saline
------------------------	---------------------------

---

<b>Preservative Stabilisers</b>	0.02% Sodium Azide (NaN <sub>3</sub> )
---------------------------------	--

---

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
---------------------------------------	----------------------------

---

<b>Immunogen</b>	An 18 amino acid peptide from near the amino terminus of human BRCC36.
------------------	--

---

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q53YX5</a> <a href="#">Related reagents</a>
--------------------------------	--

---

<b>RRID</b>	AB_2290662
-------------	------------

---

<b>Specificity</b>	<p><b>Rabbit anti Human BRCC36 antibody</b> recognizes human BRCA1/BRCA2 complex subunit 36 (BRCC36), a 36.7 kDa nuclear protein.</p> <p>BRCC36, together with BRCA1, BRCA2, Rad51 and BRCC45, forms a holoenzyme complex that possesses an ubiquitin E3 ligase activity.</p> <p>BRCC36 functions in enhancing cellular survival following DNA damage and that it is essential for ionizing radiation induced BRCA1 phosphorylation and nuclear foci formation and BRCC36 is aberrantly expressed in the majority of breast tumors indicating a potential role in the pathogenesis of the disease. BRCC36 could be a potentially important target in the treatment of radiation-resistant breast tumors (<a href="#">Chen et al. 2004</a>).</p>
--------------------	---

---

<b>Western Blotting</b>	AHP1484 detects a band of approximately 36kDa in MCF7 cell lysates.
-------------------------	---

---

<b>Further Reading</b>	<ol style="list-style-type: none"><li>1. Dong, Y. <i>et al.</i> (2003) Regulation of BRCC, a holoenzyme complex containing BRCA1 and BRCA2, by a signalosome-like subunit and its role in DNA repair. <a href="#">Mol Cell. 12 (5): 1087-99.</a></li><li>2. Chen, X. <i>et al.</i> (2006) BRCC36 is essential for ionizing radiation-induced BRCA1 phosphorylation and nuclear foci formation. <a href="#">Cancer Res. 66:5039-46.</a></li></ol>
------------------------	--

---

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
----------------	---

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP1484">https://www.bio-rad-antibodies.com/SDS/AHP1484</a>
--------------------------------------	--

---

**Regulatory**For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)Sheep Anti Rabbit IgG (STAR35...) [RPE](#)Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M416547:230302'

**Printed on 12 Aug 2023**