

Datasheet: AHP1538 BATCH NUMBER 171011

Description:	RABBIT ANTI 5-LIPOXYGENASE (pSer523)
Specificity:	5-LIPOXYGENASE (pSer523)
Other names:	5-LO
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA				
Immunoprecipitation				
Western Blotting	•			1/1000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human				
Species Cross	Reacts with: Rat				
Reactivity	Based on sequence similarity, is expected to react with:Primate				
	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or				
	personal communications from the originators. Please refer to references indicated for				
	further information.				
Product Form	Purified IgG - liquid				

Antiserum Preparation Antisera to phosphorylated 5-lipoxygenase were raised by repeated immunisations of

rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.

known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic o inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretio vasoconstriction and increased vascular permeability. I recent years 5-LO has also beer implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.		
Stabilisers 0.01% Bovine Serum Albumin 50% Glycerol Synthetic phosphopeptide corresponding to an amino acid sequence within human 5-lipoxygenase which includes the phosphorylated residue Ser523. External Database Links UniProt: P09917 Related reagents Entrez Gene: 240 ALOX5 Related reagents Synonyms LOG5 RRID AB_2226953 Specificity Rabbit anti 5-lipoxygenase (pSer523) antibody recognizes 5-lipoxygenase (5-LO), als known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic backbone 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic o inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretio vasoconstriction and increased vascular permeability. I recent years 5-LO has also beer implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.	Buffer Solution	10mM HEPES pH7.5, 150mM NaCl
External Database Links UniProt:		0.01% Bovine Serum Albumin
Links UniProt: P09917 Related reagents Entrez Gene: 240 ALOX5 Related reagents Synonyms LOG5 RRID AB_2226953 Specificity Rabbit anti 5-lipoxygenase (pSer523) antibody recognizes 5-lipoxygenase (5-LO), als known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic o inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretio vasoconstriction and increased vascular permeability. I recent years 5-LO has also beer implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.	Immunogen	
RRID Rabbit anti 5-lipoxygenase (pSer523) antibody recognizes 5-lipoxygenase (5-LO), als known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone. 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic o inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretio vasoconstriction and increased vascular permeability. I recent years 5-LO has also been implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.		P09917 Related reagents Entrez Gene:
Rabbit anti 5-lipoxygenase (pSer523) antibody recognizes 5-lipoxygenase (5-LO), als known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic or inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretion vasoconstriction and increased vascular permeability. I recent years 5-LO has also been implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.	Synonyms	LOG5
known as arachidonate 5-lipoxygenase, when phosphorylated at serine 523. 5-LO is an enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic o inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretio vasoconstriction and increased vascular permeability. I recent years 5-LO has also beer implicated in the regulation of cell proliferation and apoptosis and has been found in a broad variety of cancer cells.	RRID	AB_2226953
Western Blotting AHP1538 detects a doublet of approximately 80kDa in rat cortex lysate	Specificity	enzyme that catalyses the synthesis of leukotrienes (LTs) from arachidonic acid (a polyunsaturated fatty acid), by the insertion of an oxygen into the arachidonic backbone. 5-LO requires calcium, ATP and the 5-LO-activating protein as cofactors for full activity. Leukotrienes are synthesised and secreted by leukocytes in response to immunologic or inflammatory stimuli, such as in asthma and allergic reactions. The biological effects of LTs include leukocyte chemotaxis, pulmonary airway constriction, mucous hypersecretion, vasoconstriction and increased vascular permeability. I recent years 5-LO has also been implicated in the regulation of cell proliferation and apoptosis and has been found in a
At it 1000 detects a doublet of approximately bonds in fat cortex lysate.	Western Blotting	AHP1538 detects a doublet of approximately 80kDa in rat cortex lysate.
Storage Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this made denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.	Storage	Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
Guarantee 12 months from date of despatch	Guarantee	12 months from date of despatch
Health And Safety Information Material Safety Datasheet documentation #10088 available at: https://www.bio-rad-antibodies.com/SDS/AHP1538 10088		https://www.bio-rad-antibodies.com/SDS/AHP1538

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) FITC
Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363949:200529'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint