

Datasheet: AHP1589

BATCH NUMBER 148599

| | |
|----------------------|---------------------------------------|
| Description: | RABBIT ANTI HUMAN CX3CR1 (N-TERMINAL) |
| Specificity: | CX3CR1 (N-TERMINAL) |
| Other names: | FRACTALKINE RECEPTOR |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Immunohistology - Paraffin (1) | ▪ | | | 2.0ug/ml |
| Western Blotting | ▪ | | | 0.5 - 1.0ug/ml |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Reacts with: Rat, Mouse

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Antiserum Preparation

Antiserum to human CX3CR1 (NT) was raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

| | |
|--|---|
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.02% Sodium Azide (NaN ₃) |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml |
| Immunogen | A peptide corresponding to amino acids 2-21 of human CX3CR1. |
| External Database Links | <p>UniProt: P49238 Related reagents</p> <p>Entrez Gene: 1524 CX3CR1 Related reagents</p> |
| Synonyms | CMKBRL1, GPR13 |
| RRID | AB_2087421 |
| Specificity | <p>Rabbit anti Human CX3CR1 antibody recognizes the N-terminal region of Human CX3CR1, a 40.4 kDa seven trans-membrane protein.</p> <p>CX3CR1, also known as fractalkine receptor, is the receptor for CX3CL1 and acts as a co-receptor with CD4 for the HIV-1 envelope protein. It is expressed in a variety of human tissues and mediates leucocyte adhesion and migration.</p> |
| Histology Positive Control Tissue | Human heart |
| Western Blotting | AHP1589 detects a band of approximately 50kDa in human spleen cell lysates. |
| References | 1. Viladomiu, M. <i>et al.</i> (2017) Cooperation of Gastric Mononuclear Phagocytes with <i>Helicobacter pylori</i> during Colonization. J Immunol. 198 (8): 3195-204. |
| Further Reading | <ol style="list-style-type: none"> 1. Combadiere, C. <i>et al.</i> (1995) Cloning, chromosomal localization, and RNA expression of a human beta chemokine receptor-like gene. DNA Cell Biol. 14 (8): 673-80. 2. Combadiere, C. <i>et al.</i> (1998) Gene cloning, RNA distribution, and functional expression of mCX3CR1, a mouse chemotactic receptor for the CX3C chemokine fractalkine. Biochem Biophys Res Commun. 253 (3): 728-32. 3. Combadiere, C. <i>et al.</i> (1998) Identification of CX3CR1. A chemotactic receptor for the human CX3C chemokine fractalkine and a fusion coreceptor for HIV-1. J Biol Chem. 273 (37): 23799-804. 4. Raport, C.J. <i>et al.</i> (1995) The orphan G-protein-coupled receptor-encoding gene V28 is closely related to genes for chemokine receptors and is expressed in lymphoid and neural tissues. Gene. 163 (2): 295-9. |
| Storage | Store at +4°C or at -20°C if preferred. |

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| | |
|------------------|---------------------------------|
| Guarantee | 12 months from date of despatch |
|------------------|---------------------------------|

| | |
|--------------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1589 10040 |
|--------------------------------------|--|

| | |
|-------------------|----------------------------|
| Regulatory | For research purposes only |
|-------------------|----------------------------|

Related Products

Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Recommended Useful Reagents

- [ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)
[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M363956:200529'

Printed on 18 Jan 2024