

Datasheet: AHP1601

| Specificity:LOX-1 (EXTRACELLULAR DOMAIN)Other names:OLR1Format:SerumProduct Type:Polyclonal AntibodyIsotype:Polyclonal IgGQuantity:0.1 ml | Description: | RABBIT ANTI HUMAN LOX-1 (EXTRACELLULAR DOMAIN) |
|---|---------------|--|
| Format:SerumProduct Type:Polyclonal AntibodyIsotype:Polyclonal IgG | Specificity: | LOX-1 (EXTRACELLULAR DOMAIN) |
| Product Type: Polyclonal Antibody Isotype: Polyclonal IgG | Other names: | OLR1 |
| Isotype: Polyclonal IgG | Format: | Serum |
| | Product Type: | Polyclonal Antibody |
| Quantity: 0.1 ml | Isotype: | Polyclonal IgG |
| | Quantity: | 0.1 ml |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | |
|-----------------------------|--|-------------------|------------|-------------------------|-------------------------|--|
| | | Yes | No | Not Determined | Suggested Dilution | |
| | Flow Cytometry | | | | | |
| | Immunohistology - Frozen | | | | | |
| | Immunohistology - Paraffin | | | | | |
| | ELISA | • | | | | |
| | Immunoprecipitation | | | • | | |
| | Western Blotting | | | • | | |
| | Immunofluorescence (1) | - | | | 1/100 - 1/200 | |
| | necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) Fixation with 4% paraformaldehyde or methanol is recommended . | | | | | |
| Target Species | Human | | | | | |
| Product Form | Serum - liquid | | | | | |
| Antiserum Preparatio | n Antisera to human LOX- rabbits with highly purifie | • | | ain) were raised by re | peated immunisations of | |
| Preservative Stabilisers | 0.09% Sodium Azide (Na | aN ₃) | | | | |
| Immunogen | Peptide corresponding to | the C-ter | minal (ext | racellular) region of h | uman LOX-1. | |

| External Database Links | UniProt: P78380 Related reagents Entrez Gene: 4973 OLR1 Related reagents |
|----------------------------------|---|
| Synonyms | CLEC8A, LOX1 |
| RRID | AB_2267675 |
| Specificity | Rabbit anti Human LOX-1 antibody recognizes the extracellular (C-terminal) domain of human LOX-1, a receptor for oxidised low-density lipoproteins (LDL). This receptor mediates the recognition, internalisation and degradation of oxidised LDL. It is a Type II membrane protein with a typical C-type lectin structure at the extracellular C-terminus which recognises the ligand. LOX-1 activation by oxidised LDL causes endothelial changes such as decreased nitric oxide release and an increased expression of adhesion molecules. LOX-1 also binds activated platelets and apoptotic cells. |
| | The expression of LOX-1 is induced by proatherogenic conditions such as hyperlipidemia, hypertension and diabetes and as such appears to contribute to the pathogenesis of vascular disorders, particularly atherosclerosis (<u>Balzan & Lubrano 2018</u>). |
| References | Chen, M. <i>et al.</i> (2002) LOX-1, the receptor for oxidized low-density lipoprotein identified from endothelial cells: implications in endothelial dysfunction and atherosclerosis. <u>Pharmacol. Ther. 95: 89-100.</u> Li, D. <i>et al.</i> (2003) LOX-1, an oxidized LDL endothelial receptor, induces CD40/CD40L signaling in human coronary artery endothelial cells. <u>Arterioscler Thromb Vasc Biol. 23 (5): 816-21.</u> Morawietz, H. <i>et al.</i> (1999) Angiotensin II induces LOX-1, the human endothelial receptor for oxidized low-density lipoprotein. <u>Circulation. 100 (9): 899-902.</u> Sawamura, T. <i>et al.</i> (1997) An endothelial receptor for oxidized low-density lipoprotein. <u>Nature. 386 (6620): 73-7.</u> Yoshida, H. <i>et al.</i> (1998) Identification of the lectin-like receptor for oxidized low-density lipoprotein in human macrophages and its potential role as a scavenger receptor. <u>Biochem J. 334 (Pt 1): 9-13.</u> |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10081 available at: 10081: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10081.pdf</u> |

| Related Products | | | | | | | |
|---|--|--|--|--|--|--|--|
| Recommended Secondary Antibodies | | | | | | | |
| Sheep Anti Rabbit IgG (STAR34) FITC | | | | | | | |
| Goat Anti Rabbit IgG (H/L) (STAR124) <u>HRP</u> | | | | | | | |
| Sheep Anti Rabbit IgG (STAR35) <u>RPE</u> | | | | | | | |
| Goat Anti Rabbit IgG (Fc) (STAR121) Biotin, | <u>FITC</u> , <u>HRP</u> | | | | | | |
| Sheep Anti Rabbit IgG (STAR36) DyLigh | nt®488, DyLight®680, DyLight®800 | | | | | | |
| | | | | | | | |
| | Tel: +44 (0)1865 852 700 Europe | Tel: +49 (0) 89 8090 95 21 | | | | | |
| | Fax: +44 (0)1865 852 739 Email: antibody sales uk@bio-rad.com | Fax: +49 (0) 89 8090 95 50 Email: antibody sales de@bio-rad.com | | | | | |
| | Linali. anubouy_sales_uk@bl0-lau.com | Linali. anubody_sales_de@blo-lad.com | | | | | |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381724:210512'

Printed on 11 Jul 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint