

# Datasheet: AHP1620 BATCH NUMBER 160802

| Description:  | RABBIT ANTI HUMAN IL-27 RECEPTOR ALPHA (C-TERMINAL) |
|---------------|---|
| Specificity:  | IL-27 RECEPTOR ALPHA (C-TERMINAL)                   |
| Other names:  | TCCR  |
| Format:       | Purified  |
| Product Type: | Polyclonal Antibody                                 |
| Isotype:      | Polyclonal IgG                                      |
| Quantity:     | 0.1 mg  |

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                                | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Immunohistology - Paraffin (1) | •   |    |                | 5ug/ml             |
| Western Blotting               | -   |    |                | 0.5 - 1.0ug/ml     |
| Immunofluorescence             | -   |    |                | 20ug/ml            |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species              | Human  |
|-----------------------------|--|
| Species Cross<br>Reactivity | Reacts with: Mouse, Rat  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form                | Purified IgG - liquid  |
|                             |  |

**Antiserum Preparation** Antiserum to human IL-27Rα (C-Terminal) was raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity

chromatography.

| Buffer Solution                  | Phosphate buffered saline  |  |  |
|----------------------------------|--|--|--|
| Preservative<br>Stabilisers      | 0.02% Sodium Azide (NaN <sub>3</sub> )   |  |  |
| Approx. Protein Concentrations   | IgG concentration 0.5mg/ml   |  |  |
| Immunogen                        | A peptide corresponding to amino acids near the carboxy terminus of human IL-27R $\alpha$ precursor.   |  |  |
| External Database<br>Links       | UniProt:  Q6UWB1 Related reagents  Entrez Gene:  9466 IL27RA Related reagents  |  |  |
| Synonyms                         | CRL1, TCCR, WSX1   |  |  |
| RRID                             | AB_2295940   |  |  |
| Specificity                      | Rabbit anti human IL-27Rα antibody recognizes the carboxy terminal of the alpha subunit of the human IL-27 receptor, otherwise known as the T cell cytokine receptor (TCCR). It is a member of the type I cytokine receptor family with higher levels of expression in CD4+ T cells than in CD8+ T cells. It is a receptor for Interleukin-27 and is involved in the regulation of Th1-type T cells. It also appears to be involved in innate defense mechanisms. It is highly expressed in lymphoid tissue such as spleen, lymph nodes and peripheral blood leukocytes. It is also weakly expressed in tissues such as the heart, brain and lung. |  |  |
| Western Blotting                 | Rabbit anti human IL-27R $\alpha$ detects a band of approximately 70kDa in human spleen tissue lysates.  |  |  |
| Storage                          | Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  |  |  |
| Guarantee                        | 12 months from date of despatch  |  |  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP1620">https://www.bio-rad-antibodies.com/SDS/AHP1620</a> 10040   |  |  |
| Regulatory                       | For research purposes only   |  |  |
|                                  |  |  |  |

# **Related Products**

## **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) FITC

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

# **Recommended Useful Reagents**

#### TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363971:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint