

Datasheet: AHP1628T

BATCH NUMBER 167491

Description:	RABBIT ANTI GAPDH (N-TERMINAL)
Specificity:	GAPDH (N-TERMINAL)
Other names:	GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin (1)	▪			10ug/ml
Western Blotting	▪			0.5 - 1.0ug/ml
Immunocytochemistry	▪			1/25 - 1/100

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Mouse, Rat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Antiserum Preparation Antiserum to human GAPDH (N-Terminal) was raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity

chromatography.

Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	A peptide corresponding to a 16 amino acid sequence from near the amino-terminus of GAPDH.
External Database Links	UniProt: P04406 Related reagents Entrez Gene: 2597 GAPDH Related reagents
Synonyms	GAPD
RRID	AB_1604987
Specificity	<p>Rabbit anti GAPDH (N-Terminal) antibody recognizes human glyceraldehyde 3-phosphate dehydrogenase (GAPDH). GAPDH is a major glycolytic enzyme within the cytosol, which is also involved in a number of intracellular processes including membrane fusion, microtubule bundling, phosphotransferase, DNA replication and DNA repair.</p> <p>GAPDH may be involved in the cellular phenotype of age-related neurodegenerative disorders such as Alzheimer's and Huntington's disease.</p>
Histology Positive Control Tissue	Human liver.
Western Blotting	Rabbit anti GAPDH antibody detects a band of approximately 40kDa in HeLa cell lysates.
References	<ol style="list-style-type: none">1. Mohan, H. <i>et al.</i> (2014) Nutrients differentially regulate nucleobindin-2/nesfatin-1 <i>in vitro</i> in cultured stomach ghrelinoma (MGN3-1) cells and <i>in vivo</i> in male mice. PLoS One. 9 (12): e115102.2. Rogalska, A. <i>et al.</i> (2014) Epothilone B induces extrinsic pathway of apoptosis in human SKOV-3 ovarian cancer cells. Toxicol In Vitro. 28: 675-83.3. Paré, B. <i>et al.</i> (2015) Early detection of structural abnormalities and cytoplasmic accumulation of TDP-43 in tissue-engineered skins derived from ALS patients. Acta Neuropathol Commun. 3 (1): 5.4. Antonucci, J.M. <i>et al.</i> (2016) SAMHD1-mediated HIV-1 restriction in cells does not involve ribonuclease activity. Nat Med. 22 (10): 1072-1074.5. Nevzorova, Y.A. <i>et al.</i> (2017) Anti-tumorigenic and anti-angiogenic effects of natural conifer <i>Abies sibirica</i> terpenoids <i>in vivo</i> and <i>in vitro</i>. Biomed Pharmacother. 89: 386-95.

6. Pham, D. *et al.* (2022) Inhibition of TGFb-induced Fibroblast to Myofibroblast Transition by Conditioned Medium from Human Rhinovirus-infected Airway Epithelial Cells is Mediated by Prostaglandin E2 [Res Sq 21 Jan. \[Epub ahead of print\].](#)
7. Phillips, S. *et al.* (2022) Protocol for the generation of HIV-1 genomic RNA with altered levels of N6-methyladenosine [STAR Protocols. 3 \(3\): 101616.](#)
8. Espada, C.E. *et al.* (2023) SAMHD1 impairs type I interferon induction through the MAVS, IKKε, and IRF7 signaling axis during viral infection. [J Biol Chem. 299 \(7\): 104925.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1628T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
 Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
 Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
 Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
 'M389157:210806'

Printed on 05 Feb 2024