

Datasheet: AHP1703

BATCH NUMBER 172353

Description:	RABBIT ANTI HUMAN CD274
Specificity:	CD274
Other names:	PD-L1
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin (1)	▪			2.5ug/ml
Western Blotting	▪			0.5 - 1.0ug/ml
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Mouse, Rat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Antiserum Preparation	Antisera to Human CD274 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	A peptide corresponding to a 17 amino acid sequence from near the centre of Human CD274.
External Database Links	<p>UniProt: Q9NZQ7 Related reagents</p> <p>Entrez Gene: 29126 CD274 Related reagents</p>
Synonyms	B7H1, PDCD1L1, PDCD1LG1, PDL1
RRID	AB_2073436
Specificity	<p>Rabbit anti Human CD274 antibody detects CD274, also known as B7-H1 and PD-1L, a cell surface glycoprotein, a member of the B7 family of co-stimulatory molecules. CD274 is expressed constitutively on macrophages and dendritic cells, and is induced on activated T-cells, B-cells, endothelial cells and epithelial cells in response to Interferons alpha, beta and gamma. CD274 is reported to possess dual functions; inhibition of activated effector T cells and co-stimulation of naïve T cells. CD274 inhibits proliferation of activated T cells via ligation to the co-inhibitory molecule CD279 (programmed death-1; PD-1) leading to the secretion of the regulatory cytokine interleukin-10. CD274 has also been shown to co-stimulate early T cell priming and differentiation. Deregulated CD274 function has been reported in chronic viral and intracellular bacterial infection, as well as in many autoimmune diseases and cancers.</p>
Histology Positive Control Tissue	Human heart tissue
Western Blotting	AHP1703 detects a band of approximately 32kDa in Raji cell lysate.
References	1. Satelli, A. <i>et al.</i> (2016) Potential role of nuclear PD-L1 expression in cell-surface vimentin positive circulating tumor cells as a prognostic marker in cancer patients. Sci Rep. 6: 28910.
Further Reading	<p>1. Lagier, A.J. & Pober, J.S. (2006) Immune accessory functions of human endothelial cells are modulated by overexpression of B7-H1 (PDL1). Hum Immunol. 67 (8): 568-78.</p> <p>2. Holling, T.M. <i>et al.</i> (2004) Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. Hum Immunol. 65 (4): 282-90.</p> <p>3. Ishida, Y. <i>et al.</i> (1992) Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. EMBO J. 11 (11):</p>

[3887-95.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/AHP1703>

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M381740:210512'

Printed on 09 Dec 2025