

Datasheet: AHP1709

Description:	RABBIT ANTI HUMAN CD82 (C-TERMINAL)
Specificity:	CD82 (C-TERMINAL)
Other names:	KAI1
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Immunohistology - Paraffin (1)	•			2.5ug/ml
	Western Blotting	•			0.5 - 1.0ug/ml
	Where this product has r			•	•
	necessarily exclude its u		•		•
	a guide only. It is recomm			•	or use in their own
	system using appropriate	-	•		
	(1)This product require paraffin sections.Sodiu	-		-	
Target Species	Human				
Species Cross	Reacts with: Mouse, Rat				
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG - liquid				
Antiserum Preparation	n Antisera to Human CD82 purified antigen. Purified				• •
Buffer Solution	Phosphate buffered salin	e			

Preservative Stabilisers	0.02% Sodium Azide (NaN ₃)		
Approx. Protein Concentrations	IgG concentration 1.0mg/ml		
Immunogen	A peptide corresponding to a 15 amino acid sequence close to the carboxy terminus of Human CD82.		
External Database Links	UniProt: <u>P27701</u> <u>Related reagents</u>		
	Entrez Gene: <u>3732</u> CD82 <u>Related reagents</u>		
Synonyms	KAI1, SAR2, ST6, TSPAN27		
RRID	AB_2076397		
Specificity	 AB_2076397 Rabbit anti Human CD82 antibody recognises the carboxy-terminal region of human CD82, also known as KAI1, SAR2 and TSPAN27. CD82 is a ~30 kDa multi-pass transmembrane glycoprotein belonging to the tetraspanin (TM4SF) family. CD82 acts as a cancer metastasis suppressor protein (Feng <i>et al.</i>2015), while ubiquitously expressed at moderate to high levels in many normal tissues, its expression is down regulated during tumour progression. Loss of CD82 and p53 is correlated with poor prognosis for patients with prostate and other forms of cancer (Liu <i>et al.</i> 2006). CD82 interacts with CD234 (DARC). Tumour cells dislodged from the primary tumour and expressing CD82 interact with CD234 expressed on vascular cells, transmitting a senescent signal to the tumour cells (Tsai <i>et al.</i>2011). Tumor cells that have lost CD82 expression can proliferate and potentially give rise to metastases (Miranti 2009). Two isoforms of CD82 may be generated by alternative splicing. The c-terminal region is not affected and thus Rabbit anti Human CD82 antibody is expected to recognize both isoforms of human CD82 (UniProt: P27701). 		
Histology Positive Control Tissue	Human colon tissue		
Western Blotting	AHP1709 detects a band of approximately 33kDa in A549 cell lysate		
References	1. Zuidscherwoude, M. <i>et al.</i> (2015) The tetraspanin web revisited by super-resolution microscopy. <u>Sci Rep. 5: 12201.</u>		
Further Reading	1. Kauffman, E.C. <i>et al.</i> (2003) Metastasis suppression: the evolving role of metastasis suppressor genes for regulating cancer cell growth at the secondary site. <u>J Urol. 169 (3):</u> <u>1122-33.</u>		

	 Dong, J.T. <i>et al.</i> (1995) KAI1, a metastasis suppressor generic human chromosome 11p11.2. <u>Science. 268 (5212): 884-6.</u> Gil, M.L. <i>et al.</i> (1992) A member of the tetra spans transmeric is recognized by a monoclonal antibody raised against an HLA lymphokine-activated killer-susceptible, B lymphocyte line. Clofunctional studies. <u>J Immunol. 148 (9): 2826-33.</u> Gaugitsch, H.W. <i>et al.</i> (1991) A new superfamily of lymphoid proteins with extensive homology to <i>Schistosoma mansoni</i> and <u>21 (2): 377-83.</u> 	mbrane protein superfamily class I-deficient, ning and preliminary d and melanoma cell	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP1709 10040		
Regulatory	For research purposes only		

Related Products

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M416556:230302'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint