

Datasheet: AHP1734

**BATCH NUMBER 161212**

<b>Description:</b>	RABBIT ANTI HUMAN TGF BETA
<b>Specificity:</b>	TGF BETA
<b>Other names:</b>	TRANSFORMING GROWTH FACTOR BETA
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin (1)	▪			5ug/ml
ELISA	▪			1/2000 - 1/10000
Western Blotting	▪			1/200 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Rabbit, Mouse, Rat</p> <p>Based on sequence similarity, is expected to react with:Mammals</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid

**Antiserum Preparation** Antisera to Human TGF-beta1 were raised by repeated immunisation of Rabbits with

highly purified antigen. Purified IgG prepared from whole serum by affinity chromatography.

---

**Buffer Solution** Phosphate buffered saline

---

**Preservative Stabilisers** 0.09% Sodium Azide (NaN<sub>3</sub>)

---

**Approx. Protein Concentrations** IgG concentration 0.4 mg/ml.

---

**Immunogen** A synthetic peptide corresponding to amino acids 286-293 of the human TGF-beta precursor protein.

---

**External Database Links**

**UniProt:**

[P01137](#) [Related reagents](#)

**Entrez Gene:**

[7040](#) TGFB1 [Related reagents](#)

---

**Synonyms** TGFB

---

**RRID** AB\_2222443

---

**Specificity** **Rabbit anti Human TGF beta antibody** recognizes human Transforming Growth Factor Beta 1 (TGF-beta1), a 390 amino acid secreted protein belonging to the TGF-beta family.

TGF-beta1 is a multifunctional protein that controls proliferation, differentiation and other functions in a wide range of cell types and both positively and negatively regulates many other growth factors. TGF-beta1 also plays an important role in bone remodelling through stimulation of osteoblastic bone formation causing chemotaxis, proliferation and differentiation in committed osteoblasts.

Defects in TGFB1 are the cause of Camurati-Engelmann disease (CED), also known as progressive diaphyseal dysplasia 1 ([Kinoshita et al. 2000](#)).

---

**Histology Positive Control Tissue** Human spleen

---

**References**

1. Polichnowski, A.J. *et al.* (2014) Severe renal mass reduction impairs recovery and promotes fibrosis after AKI. [J Am Soc Nephrol. 25 \(7\): 1496-507.](#)
2. Moore, L.B. *et al.* (2015) Loss of monocyte chemoattractant protein-1 alters macrophage polarization and reduces NFκB activation in the foreign body response. [Acta Biomater. 11: 37-47.](#)
3. Geng, H. *et al.* (2012) Lysophosphatidic acid increases proximal tubule cell secretion of profibrotic cytokines PDGF-B and CTGF through LPA2- and Gαq-mediated Rho and αvβ6 integrin-dependent activation of TGF-β. [Am J Pathol. 181: 1236-49.](#)
4. Tark, K.C. *et al.* (2015) Effects of ginsenoside Rb1 on hypertrophic scar remodeling in

rabbit model. [Eur J Pharmacol. 750C: 151-59.](#)

---

**Storage** Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10094 available at:  
<https://www.bio-rad-antibodies.com/SDS/AHP1734>  
10094

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M364011:200529'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)