Datasheet: AHP1776T BATCH NUMBER 240414

Description:	RABBIT ANTI RIG-I
Specificity:	RIG-I
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications	ns This product has been reported to work in the following applications. This inform derived from testing within our laboratories, peer-reviewed publications or person communications from the originators. Please refer to references indicated for ful information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
	Suggested Dilution						
	Immunohistology - Paraffin (1)	-			5.0ug/ml		
	Western Blotting	-			0.5 - 1.0ug/ml		
	Where this product has not been tested for use in a particular technique this does not						
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.						
Target Species	Human						
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity a reactivity is derived from personal communications further information.	testing w	ithin our la	aboratories, peer-revie	wed publications or		
Product Form	Purified IgG - liquid						
Antiserum Preparatio	on Antisera to human RIG-I purified antigen. Purified						
Buffer Solution	Phosphate buffered salin	e					

Preservative Stabilisers	0.02% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Human GST-tagged RIG-1 protein.
External Database Links	UniProt: 095786 Related reagents
	Entrez Gene: <u>23586</u> DDX58 <u>Related reagents</u>
RRID	AB_2175703
Specificity	Rabbit anti RIG-I antibody recognizes human Retinoic acid-inducible gene 1 protein (RIG-I), also known as RIG-1 or DEAD-box protein 58. RIG-I is a receptor found in the cytoplasm that recognizes viral RNA from replicating viruses in infected cells. Upon activation with intracellular RNA, a cascade is triggered resulting in the activation of NF-kappa-B, and the eventual induction of antiviral cytokines such as IFN-beta and RANTES. RIG-I is essential for the production of interferons in response to RNA viruses and as such play a key role in innate immunity.
Histology Positive Control Tissue	Human heart
	Human heart AHP1776T detects a band of approximately 84 kDa in C2C12cell lysate.
Control Tissue	
Control Tissue Western Blotting	 AHP1776T detects a band of approximately 84 kDa in C2C12cell lysate. 1. Satoh, T. <i>et al.</i> (2014) Tumor microenvironment and RIG-I signaling molecules in Epstein Barr virus-positive and -negative classical Hodgkin lymphoma of the elderly. <u>J Clin</u>
Control Tissue Western Blotting References	 AHP1776T detects a band of approximately 84 kDa in C2C12cell lysate. 1. Satoh, T. <i>et al.</i> (2014) Tumor microenvironment and RIG-I signaling molecules in Epstein Barr virus-positive and -negative classical Hodgkin lymphoma of the elderly. J Clin Exp Hematop. 54 (1): 75-84. 1. Akira, S. <i>et al.</i> (2006) Pathogen recognition and innate immunity. Cell. 124 (4): 783-801. 2. Yoneyama, M. <i>et al.</i> (2004) The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. Nat Immunol. 5 (7): 730-7. 3. Sharma S, et al (2003) Triggering the interferon antiviral response through an IKK-related pathway Science 300:1148-51. 4. Alexopoulou, L. <i>et al.</i> (2001) Recognition of double-stranded RNA and activation of

Health Ar Informati		://www.bio-rad-ar	heet documentation #10 htibodies.com/SDS/AHP		le at:
Regulato	ry For r	esearch purpose	s only		
Relate	d Products				
Recomn	nended Useful R	eagents			
ANTIGEN	RETRIEVAL BUFFE	R, pH8.0 (BUF025	<u>A)</u>		
North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751 Email: antibody_sales_us	@bio-rad.com	Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	o-rad.com	Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specific data	sheet for this produ	uct, please use our online 'M364033:200529'	search tool at	: bio-rad-antibodies.com/datasheets
			Printed on 18 Jan 2024		

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint