

Datasheet: AHP1784T

**BATCH NUMBER 161017**

<b>Description:</b>	RABBIT ANTI SIGIRR (C-TERMINAL)
<b>Specificity:</b>	SIGIRR (C-TERMINAL)
<b>Other names:</b>	TIR8
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			10ug/ml
Western Blotting	▪			0.5 - 1ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

### Target Species

Mouse

### Species Cross Reactivity

Reacts with: Human

Based on sequence similarity, is expected to react with:Rat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG - liquid

### Antiserum Preparation

Antiserum to SIGIRR (CT) was raised by repeated immunisation of with highly purified antigen. Purified IgG was prepared from whole serum by immunoaffinity chromatography.

### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.02% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	A synthetic peptide corresponding to 15 amino acids near the carboxy terminus of mouse SIGIRR.
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">Q9JLZ8</a>    <a href="#">Related reagents</a></p> <p><a href="#">Q6IA17</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">24058</a>    Sigirr    <a href="#">Related reagents</a></p> <p><a href="#">59307</a>    SIGIRR    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Tir8
<b>RRID</b>	AB_2188896
<b>Specificity</b>	<p><b>Rabbit anti SIGIRR (C-terminal) antibody</b> recognizes the Single Ig IL-1-related receptor, also known as SIGIRR, Single immunoglobulin domain-containing IL1R-related protein, Toll/interleukin-1 receptor 8 or TIR8. SIGIRR is a 409 amino acid ~ 45-90 kDa single pass type III transmembrane glycoprotein, Multiple species of SIGIRR seen in western blotting are achieved through differential glycosylation at four potential glycosylation sites in the SIGIRR ectodomain, mutation at these sites to eliminate glycosylation yields a single band of the predicted size (44 kDa) for unmodified SIGIRR (<a href="#">Zhao et al. 2015</a>).</p> <p>SIGIRR is a member of the Toll-like receptor-interleukin 1 receptor superfamily, defined by an intracellular Toll-IL-1R (TIR) domain. SIGIRR acts as a negative regulator of the Toll-like and IL-1R receptor signalling pathways. It interacts with TLR4, TLR5, TLR9 and IL-1R through the extracellular Ig domain, or the intracellular TIR domain. The Ig domain interferes with the heterodimerization of IL-1R while the intracellular TIR domain inhibits both IL-1 and LPS signalling by attenuating the recruitment of receptor-proximal signalling components to TLR4. SIGIRR is expressed in kidney, colon, small intestine, lung, spleen and liver. It is not expressed in brain or muscle.</p> <p>Rabbit anti SIGIRR antibody has been successfully used for the detection of SIGIRR in both murine and human samples by western blotting (<a href="#">Zhao et al. 2015</a>)</p>
<b>Western Blotting</b>	AHP1784T detects a band of approximately 43kDa in A549 cell lysate
<b>References</b>	1. Zhao, J. <i>et al.</i> (2015) Human Colon Tumors Express a Dominant-Negative Form of SIGIRR That Promotes Inflammation and Colitis-Associated Colon Cancer in Mice. <a href="#">Gastroenterology. 149 (7): 1860-1871.e8.</a>

**Storage** Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/AHP1784T>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)  
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)  
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)  
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M364039:200529'

Printed on 12 Aug 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)