Datasheet: AHP1797 BATCH NUMBER 158735

Description:	RABBIT ANTI RIP3
Specificity:	RIP3
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	plications This product has been reported to work in the following applications. This information				
derived from testing within our laboratories, peer-reviewed publications or p					ations or personal
	communications from the	originator	rs. Please	refer to references ir	ndicated for further
	information. For general	protocol re	commend	ations, please visit <u>v</u>	www.bio-
	rad-antibodies.com/proto	cols.			
		Yes	No	Not Determined	Suggested Dilution
	Immunohistology - Paraffin (1)				5ug/ml
	Western Blotting				0.5 - 1.0ug/ml
	Where this product has n necessarily exclude its us a guide only. It is recomm system using appropriate (1) This product require	ot been te se in such nended tha negative/ s antigen	ested for us procedure at the user positive cc retrieval u	se in a particular tech s. Suggested workin titrates the product to ontrols. using heat treatmer	hnique this does not ng dilutions are given as for use in their own n t prior to staining of
	paraffin sections.Sodiu	m citrate	buffer pH	6.0 is recommende	ed for this purpose.
Target Species	paraffin sections.Sodiu Mouse	m citrate	buffer pH	6.0 is recommende	ed for this purpose.
Target Species Species Cross	paraffin sections.Sodiu Mouse Reacts with: Rat	m citrate	buffer pH	6.0 is recommende	ed for this purpose.
Target Species Species Cross Reactivity	paraffin sections.Sodiu Mouse Reacts with: Rat N.B. Antibody reactivity a reactivity is derived from personal communications further information.	m citrate and workin testing wit s from the	buffer pH g conditior hin our lab originators	6.0 is recommende	ed for this purpose. In species. Cross ewed publications or erences indicated for
Target Species Species Cross Reactivity Product Form	paraffin sections.SodiuMouseReacts with: RatN.B. Antibody reactivity areactivity is derived frompersonal communicationsfurther information.Purified IgG - liquid	m citrate and workin testing wit s from the	buffer pH g conditior hin our lab originators	6.0 is recommende	ed for this purpose. In species. Cross ewed publications or erences indicated for
Target Species Species Cross Reactivity Product Form Antiserum Preparation	paraffin sections.Sodiu Mouse Reacts with: Rat N.B. Antibody reactivity a reactivity is derived from personal communications further information. Purified IgG - liquid Antisera to mouse RIP3 v purified antigen. Purified	m citrate and workin testing wit s from the were raise IgG prepa	buffer pH g conditior hin our lab originators d by repea red from w	6.0 is recommended his may vary betweer poratories, peer-revie s. Please refer to refer to refer to refer to refer to refer to refer	ed for this purpose. In species. Cross ewed publications or erences indicated for
Target Species Species Cross Reactivity Product Form Antiserum Preparation Buffer Solution	paraffin sections.Sodiu Mouse Reacts with: Rat N.B. Antibody reactivity a reactivity is derived from personal communications further information. Purified IgG - liquid Antisera to mouse RIP3 of purified antigen. Purified Phosphate buffered salin	m citrate and workin testing wit s from the were raise IgG prepa e	buffer pH g conditior hin our lab originators d by repea red from w	6.0 is recommended hs may vary betweer poratories, peer-revie s. Please refer to refer ated immunisation of hole serum by affinit	ed for this purpose.

Preservative Stabilisers	0.02% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Synthetic peptide sequence corresponding to amino acids 473-486 of mouse RIP3.
External Database Links	UniProt: Q9QZL0 Related reagents Entrez Gene: 56532 Ripk3 Related reagents
Synonyms	Rip3
RRID	AB_2178676
Specificity	Rabbit anti RIP3 antibody recognizes mouse serine-threonine kinase receptor-interacting protein 3 (RIP3). RIP3 is related to RIP and RIP2 but does not posses a death domain or CARD motif at its C-terminus. RIP3 activates NF-Kappa B and potently induces apoptosis. RIP3 is expressed in a range of tissues including the spleen, liver, testis, heart, brain and lung.
Histology Positive Control Tissue	Rat kidney
Western Blotting	AHP1797 detects a band of approximately 56 kDa in Mouse 3T3 whole cell lysate
References	 Feng, S. <i>et al.</i> (2007) Cleavage of RIP3 inactivates its caspase-independent apoptosis pathway by removal of kinase domain. <u>Cell Signal. 19 (10): 2056-67.</u> Günther, C. <i>et al.</i> (2011) Caspase-8 regulates TNF-α-induced epithelial necroptosis and terminal ileitis. <u>Nature. 477: 335-9.</u> Mizumura, K. <i>et al.</i> (2014) Mitophagy-dependent necroptosis contributes to the pathogenesis of COPD. J Clin Invest. 124 (9): 3987-4003. Huang, S.C. <i>et al.</i> (2014) Tumor necrosis factor suppresses NR5A2 activity and intestinal glucocorticoid synthesis to sustain chronic colitis. <u>Sci Signal. 7 (314): ra20.</u> Meng, L. <i>et al.</i> (2015) RIP3-mediated necrotic cell death accelerates systematic inflammation and mortality. <u>Proc Natl Acad Sci U S A. 112 (35): 11007-12.</u> Mizumura, K. <i>et al.</i> (2018) Sphingolipid regulation of lung epithelial cell mitophagy and necroptosis during cigarette smoke exposure. <u>FASEB J. 32 (4): 1880-90.</u> Najafov, A. <i>et al.</i> (2019) TAM Kinases Promote Necroptosis by Regulating Oligomerization of MLKL. <u>Mol Cell. 75 (3): 457-468.e4.</u> Matsuzawa-Ishimoto, Y. <i>et al.</i> (2022) The γδ IEL effector API5 masks genetic susceptibility to Paneth cell death. <u>Nature. 610 (7932): 547-54.</u>

Further Reading 1. Yu, P.W. *et al.* (1999) Identification of RIP3, a RIP-like kinase that activates apoptosis

	and NFkappaB. <u>Curr Biol. 9 (10): 539-42.</u>			
	2. Sun, X., et al (1999) RIP3, a novel apoptosis-inducing kinase. <u>J Biol Chem.</u>			
	<u>274:16871-5.</u>			
	3. Pazdernik, N.J., et al (1999) Mouse receptor interacting prot	tein 3 does not contain a		
	caspase-recruiting or a death domain but induces apoptosis ar	nd activates NF-κB. <u>Mol Cell</u>		
	<u>Bio.19:6500-8</u>			
Storage	This product is shipped at ambient temperature. It is recomme	nded to aliquot and store at		
	-20°C on receipt. When thawed, aliquot the sample as needed	. Keep aliquots at 2-8°C for		
	short term use (up to 4 weeks) and store the remaining aliquot	s at -20°C.		
	Avoid repeated freezing and thawing as this may denature the	antibody. Storage in		
	frost-free freezers is not recommended.			
Guarantee	12 months from date of despatch			
Health And Safety	Material Safety Datasheet documentation #10040 available at:			
Information	https://www.bio-rad-antibodies.com/SDS/AHP1797			
	10040			
Regulatory	For research purposes only			

Related Products

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376 Worldw	wide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382827:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint