

Datasheet: AHP1821 BATCH NUMBER 180319

| Description: | GOAT ANTI HUMAN INTERLEUKIN-23 RECEPTOR (N-TERMINAL) |
|---------------|--|
| Specificity: | IL-23 RECEPTOR (N-TERMINAL) |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.1 mg |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | | |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin (1) | • | | | 1/125 |
| ELISA | | | | |
| Immunoprecipitation | | | | |
| Western Blotting | • | | | 1/500 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections

| Target Species | Human |
|-----------------------------|---|
| Species Cross Reactivity | Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form | Purified IgG - liquid |

Antiserum Preparation Antiserum to human IL-23R (NT) was raised by repeated immunisation of goats with

highly purified antigen. Purified IgG was prepared by affinity chromatography.

| Buffer Solution | Phosphate buffered saline | | | |
|--------------------------------------|---|--|--|--|
| Preservative | 0.1% Sodium Azide (NaN ₃) | | | |
| Stabilisers | 0.1% Bovine Serum Albumin | | | |
| | | | | |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml | | | |
| Immunogen | Synthetic peptide CQPRKLHFYKNGIKERFQITRINKTTARL corresponding to amino acids 59-87 within the N-terminal region of human IL-23 receptor. | | | |
| External Database | | | | |
| Links | UniProt: | | | |
| | Q5VWK5 Related reagents | | | |
| | Q8K4B4 Related reagents | | | |
| | Entrez Gene: | | | |
| | | | | |
| | 149233 IL23R Related reagents 209590 II23r Related reagents | | | |
| | 209590 II23r Related reagents | | | |
| RRID | AB_2233639 | | | |
| Specificity | Goat anti Human Interleukin-23 receptor antibody recognizes the human interleukin-23 receptor (IL-23R). The interleukin-23 receptor is 629 amino acid ~72kDa single pass type I transmembrane glycoprotein belonging to the type I cytokine receptor family and possessing two fibronectin type III domains. Multiple potential isoforms formed by alternative splicing are described. Goat anti Human Interleukin-23 receptor antibody recognizes an epitope within the N-terminal (NT) region of Interleukin-23 receptor (IL-23R), present in isoforms 1, 3 and 4 (UniProt: Q5VWK5), the antibody is not expected to recognize other truncated forms of the receptor lacking the immunogen sequence. IL-23R is a mediator of T-cell, NK cell, and possibly macrophage/myeloid cell, stimulation acting as the receptor for IL-23, by association with the beta component of IL-12R and is thought to signal through activation of the JAK-STAT signaling pathway. Genetic variation in IL23R is associated with inflammatory bowel disease type 17 (Kistner et al. 2006) and susceptibility to the ulcerative and inflammatory psoriasis (Hüffmeier et al. 2009). Goat anti Human Interleukin-23 receptor antibody is reported as suitable for use in immunocytochemistry on acetone fixed cells and has been used successfully for the immunohistochemical localization of IL-23R in formalin fixed, paraffin embedded human tissues. | | | |
| Histology Positive Control Tissue | Spleen | | | |
| Storage | Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. | | | |

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/AHP1821 10041 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364058:200529'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint