

Datasheet: AHP1822 BATCH NUMBER 146995

Description:	GOAT ANTI HUMAN CD284 (N-TERMINAL)
Specificity:	CD284 (N-TERMINAL)
Other names:	TLR4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit					

Antiserum Preparation Antiserum to human CD284 (NT) was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline			
Preservative	0.1% Sodium Azide (NaN ₃)			
Stabilisers	0.1% Bovine Serum Albumin			
Approx. Protein Concentrations	IgG concentration 1.0mg/ml			
Immunogen	Synthetic peptide LIQSFKLPEYFSNLTNLEHLDLSSNKIQSIYC corresponding to amino acids 161-192 within the N-terminal region of Human CD284.			
External Database	UniProt:			
Links				
	O00206 Related reagents			
	Entrez Gene:			
	7099 TLR4 Related reagents			
RRID				
KKID	AB_2256158			
Specificity	Goat anti Human CD284 antibody recognizes the human Toll-like receptor 4, also known			
	as CD284, hToll or TLR4. CD284 is an 839 amino acid ~96 kDa single pass type I			
	transmembrane glycoprotein expressed by peripheral blood leucocytes, activated T cells,			
	monocytes, macrophages, dendritic cells, granulocytes and endothelial cells. CD284 plays			
	a primary role in the activation of innate immunity (<u>UniProt: 000206</u>).			
	Goat anti Human CD284 antibody recognizes an epitope within the N-terminal (NT) region			
	of CD284, a highly conserved member of the toll-like receptor family which acts as a			
	receptor for bacterial lipopolysaccharides (LPS) in co-operation with CD14 and MD2			
	(LY96), resulting in the activation of the MyD88-dependent and MyD88-independent			
	(TRIF-dependent) signalling pathways and the production of inflammatory cytokines.			
	LPS-activated CD284 can also induce dendritic cell maturation via TRIF-dependent			
	signaling.			
	ograng.			
	Goat anti Human CD284 antibody is reported to be suitable for use in			
	immunocytochemistry on acetone fixed cells.			
Histology Positive	Calcon entensil			
Control Tissue	Spleen or tonsil			
Storage	Store at +4°C or at -20°C if preferred.			
	Storage in frost-free freezers is not recommended.			
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may			
	denature the antibody. Should this product contain a precipitate we recommend			
	microcentrifugation before use.			

Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/AHP1822 10041				
Regulatory	For research purposes only				
Related Products					
Recommended Secondary Antibodies					

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & South	Tel: +1 800 265 7376	Wor
America	Fax: +1 919 878 3751	
	Email: antibody_sales_	us@bio-rad.com

rldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364059:200529'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint