

# Datasheet: AHP1918 BATCH NUMBER 156788

| Description:  | RABBIT ANTI HUMAN PLATELET FACTOR-4 |
|---------------|-------------------------------------|
| Specificity:  | PLATELET FACTOR-4                   |
| Other names:  | CXCL4                               |
| Format:       | Purified                            |
| Product Type: | Polyclonal Antibody                 |
| Isotype:      | Polyclonal IgG                      |
| Quantity:     | 0.1 mg                              |
|               |                                     |

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                                | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry                 |     |    |                |                    |
| Immunohistology - Frozen       |     |    |                |                    |
| Immunohistology - Paraffin (1) | •   |    |                |                    |
| ELISA                          |     |    |                | 0.5 - 2.0ug/ml     |
| Immunoprecipitation            |     |    |                |                    |
| Western Blotting               |     |    |                | 0.1 - 0.2ug/ml     |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species | Human   |
|----------------|---|
| Product Form   | Purified IgG - lyophilised  |
| Reconstitution | Reconstitute with 0.1ml distilled water  Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.  For long term storage the addition of 0.09% sodium azide is recommended.  N.B. For functional studies do not add sodium azide |

**Antiserum Preparation** Antiserum to human PF-4 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

| Buffer Solution                   | Phosphate buffered saline   |
|-----------------------------------|---|
| Preservative<br>Stabilisers       | None present  |
| Carrier Free                      | Yes   |
| Approx. Protein<br>Concentrations | IgG concentration 1.0mg/ml after reconstitution   |
| Immunogen                         | Recombinant human PF-4.   |
| External Database<br>Links        | UniProt: P02776 Related reagents  Entrez Gene: 5196 PF4 Related reagents  |
| Synonyms                          | CXCL4, SCYB4  |
| RRID                              | AB_2299453  |
| Specificity                       | Rabbit anti Human Platelet Factor-4 antibody recognizes platelet factor-4 (PF-4), otherwise known as CXCL4, a member of the intercrine alpha (chemokine CXC) family, released from the alpha granules of activated platelets, during platelet aggregation.  PF-4 binds with high affinity to heparin, counteracting its' anticoagulant effect, thereby promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 acts as a chemotactic factor for neutrophils, monocytes and fibroblasts, and binds wih CXCR3B, a splice variant of the chemokine receptor CXCR3. |
| Storage                           | Prior to reconstitution store at -20°C.  After reconstitution store at -20°C.  This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  |
| Guarantee                         | 12 months from date of despatch   |
| Health And Safety<br>Information  | Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP1918">https://www.bio-rad-antibodies.com/SDS/AHP1918</a> 10162  |
| Regulatory                        | For research purposes only  |
|                                   |   |

# **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

## **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372512:200706'

#### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint