

Datasheet: AHP1918

Description:	RABBIT ANTI HUMAN PLATELET FACTOR-4
Specificity:	PLATELET FACTOR-4
Other names:	CXCL4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

#### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	<b>Suggested Dilution</b>
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)	•			
ELISA				0.5 - 2.0ug/ml
Immunoprecipitation			•	
Western Blotting	•			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Human
Purified IgG - lyophilized
Reconstitute with 0.1ml distilled water
Care should be taken during reconstitution as the protein may appear as a film at the
bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.
For long term storage the addition of 0.09% sodium azide is recommended.
N.B. For functional studies do not add sodium azide

Antiserum Preparation Antiserum to human PF-4 were raised by repeated immunisation of rabbits with highly

purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution		
	Phosphate buffered saline	
Preservative Stabilisers	None present	
Carrier Free	Yes	
Approx. Protein Concentrations	IgG concentration 1.0mg/ml after reconstitution	
Immunogen	Recombinant human PF-4.	
External Database Links	UniProt: P02776 Related reagents  Entrez Gene: 5196 PF4 Related reagents	
Synonyms	CXCL4, SCYB4	
RRID	AB_2299453	
Specificity	Rabbit anti Human Platelet Factor-4 antibody recognizes platelet factor-therwise known as CXCL4, a member of the intercrine alpha (chemoking released from the alpha granules of activated platelets, during platelet a	ne CXC) family,
	PF-4 binds with high affinity to heparin, counteracting its' anticoagulant of promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 chemotactic factor for neutrophils, monocytes and fibroblasts, and binds splice variant of the chemokine receptor CXCR3.	effect, thereby 4 acts as a
Storage	promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 chemotactic factor for neutrophils, monocytes and fibroblasts, and binds	effect, thereby 4 acts as a
Storage	promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 chemotactic factor for neutrophils, monocytes and fibroblasts, and binds splice variant of the chemokine receptor CXCR3.  This product is shipped at ambient temperature.  Prior to reconstitution store at -20°C.	effect, thereby 4 acts as a 5 wih CXCR3B, a 6 not 1 ire the antibody.
Storage	promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 chemotactic factor for neutrophils, monocytes and fibroblasts, and binds splice variant of the chemokine receptor CXCR3.  This product is shipped at ambient temperature. Prior to reconstitution store at -20°C. After reconstitution store at -20°C.  This product should be stored undiluted. Storage in frost-free freezers is recommended. Avoid repeated freezing and thawing as this may denated	effect, thereby 4 acts as a 5 wih CXCR3B, a 6 not 1 ire the antibody.
	promoting blood coagulation, and implicating PF-4 in wound repair. PF-4 chemotactic factor for neutrophils, monocytes and fibroblasts, and binds splice variant of the chemokine receptor CXCR3.  This product is shipped at ambient temperature. Prior to reconstitution store at -20°C. After reconstitution store at -20°C.  This product should be stored undiluted. Storage in frost-free freezers is recommended. Avoid repeated freezing and thawing as this may denature. Should this product contain a precipitate we recommend microcentrifugation.	effect, thereby 4 acts as a 5 wih CXCR3B, a 6 not 1 ire the antibody.

# **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

## **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M439036:250523'

#### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint