

Datasheet: AHP1919B

Description:	RABBIT ANTI HUMAN CCL17:Biotin
Specificity:	CCL17
Other names:	TARC
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.25 - 1.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - lyophilised
Reconstitution	Reconstitute with 0.5ml sterile PBS containing 0.1% bovine serum albumin Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended.
Antiserum Preparation	Antiserum to human CCL17 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Approx. Protein Concentrations	IgG concentration 0.1mg/ml after reconstitution

Immunogen Recombinant human CCL17

External Database Links

UniProt:
[Q92583](#) [Related reagents](#)

Entrez Gene:
[6361](#) CCL17 [Related reagents](#)

Synonyms SCYA17, TARC

Specificity **Rabbit anti Human CCL17 antibody** specifically recognizes human CCL17, otherwise known as TARC (thymus and activation-regulated chemokine), a member of the intercrine beta (chemokine CC) family, predominantly produced by thymic dendritic cells, which specifically acts as a chemotactic factor for T cells, and signals through binding to the CCR4 receptor.

ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with [AHP1919](#) as the capture antibody.

Storage Prior to reconstitution store at +4°C.
After reconstitution store at -20°C.
Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 12 months from date of reconstitution.

Health And Safety Information Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M320657:180726'

Printed on 18 Sep 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)