

## Datasheet: AHP1919B

<b>Description:</b>	RABBIT ANTI HUMAN CCL17:Biotin
<b>Specificity:</b>	CCL17
<b>Other names:</b>	TARC
<b>Format:</b>	Biotin
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.25 - 1.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG conjugated to Biotin - lyophilised
<b>Reconstitution</b>	Reconstitute with 0.5ml sterile PBS containing 0.1% bovine serum albumin Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended.
<b>Antiserum Preparation</b>	Antiserum to human CCL17 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1mg/ml after reconstitution

**Immunogen** Recombinant human CCL17.

---

**External Database Links**

**UniProt:**  
[Q92583](#)   [Related reagents](#)

**Entrez Gene:**  
[6361](#) CCL17   [Related reagents](#)

---

**Synonyms** SCYA17, TARC

---

**Specificity** **Rabbit anti Human CCL17 antibody** specifically recognizes human CCL17, otherwise known as TARC (thymus and activation-regulated chemokine), a member of the intercrine beta (chemokine CC) family, predominantly produced by thymic dendritic cells, which specifically acts as a chemotactic factor for T cells, and signals through binding to the CCR4 receptor.

---

**ELISA** This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with [AHP1919](#) as the capture antibody.

---

**Storage** Prior to reconstitution store at +4°C.  
After reconstitution store at -20°C.  
Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Shelf Life** 12 months from date of reconstitution.

---

**Health And Safety Information** Material Safety Datasheet documentation #10162 available at:  
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

---

**Regulatory** For research purposes only

---

**North & South America** Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M332329:181003'

**Printed on 06 Dec 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)