

## Datasheet: AHP2116 BATCH NUMBER 158492

Description:	GOAT ANTI HUMAN DESMIN (C-TERMINAL)
Specificity:	DESMIN (C-TERMINAL)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

## **Product Details**

	communications from the information. For general rad-antibodies.com/proto Flow Cytometry Immunohistology - Frozen Immunohistology - Paraffin (1) ELISA	protocol re				
	rad-antibodies.com/proto Flow Cytometry Immunohistology - Frozen Immunohistology - Paraffin (1)	<u>cols</u> . Yes		Not Determined		
	Flow Cytometry Immunohistology - Frozen Immunohistology - Paraffin (1)	Yes	No	•	Suggested Dilution	
	Immunohistology - Frozen Immunohistology - Paraffin (1)		Νο	•	Suggested Dilution	
	Immunohistology - Frozen Immunohistology - Paraffin (1)			•		
	Immunohistology - Paraffin (1)	•		-		
	(1)	-				
					3.0 - 6.0ug/ml	
		-			1/2000	
	Immunoprecipitation					
	Western Blotting				0.1 - 0.3ug/ml	
	<ul> <li>Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.</li> <li>(1) This product requires heat-mediated antigen retrieval prior to staining of paraffin sections. Sodium citrate buffer pH6.0 is recommended for this purpose.</li> </ul>					
Target Species	Human					
	Reacts with: Mouse, Rat, Pig Based on sequence similarity, is expected to react with:Dog, Bovine <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG - liquid					

Antiserum Preparation Antiserum to human Desmin (CT) was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	TRIS buffered saline				
Preservative	0.02% Sodium Azide (NaN <sub>3</sub> )				
Stabilisers	0.5% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml				
Immunogen	Synthetic peptide sequence C-RDGEVVSEATQQQHE from the C-Terminal region of Desmin (NP_001918.3).				
External Database					
Links	UniProt:				
	P17661 Related reagents				
	Entrez Gene:				
	1674 DES Related reagents				
RRID	AB_2092438				
Specificity	<b>Goat anti Human Desmin antibody</b> recognizes an epitope w region of human Desmin, a type III intermediate filament (IF), sacromeres of skeletal, smooth and cardiac muscle tissue.				
	Desmin plays an essential role in the architecture of muscle ce the subsarcolemmal cytoskeleton, and the contractile apparate nucleus and post-synaptic areas of motor endplates. Desmin i muscle tissue in embryogenesis, and a marker for tumours of in the Desmin gene, are responsible for the neuromuscular dis related cardio skeletal myopathy (CSM), characterized by skel	us to the, mitochondria, cell is an early indicator for myogenic origin. Mutations sorder known as desmin-			
Histology Positive Control Tissue	Human prostate				
Western Blotting	<b>Goat anti Human Desmin</b> detects a band of approximately 55 muscle cell lysates.	5kDa in human skeletal			
Storage	This product is shipped at ambient temperature. It is recomme -20°C on receipt. When thawed, aliquot the sample as needed short term use (up to 4 weeks) and store the remaining aliquo	I. Keep aliquots at 2-8°C for			
	Avoid repeated freezing and thawing as this may denature the frost-free freezers is not recommended.	e antibody. Storage in			
Guarantee	12 months from date of despatch				

Health A Informati		at:			
Regulato	ry	For research purpose			
Relate	d Products	S			
Recomm	nended Sec	ondary Antibodies	6		
Rabbit A	nti Goat IgG (I	Fc) (STAR122) <u>FITC</u>	<u>), HRP</u>		
Recomm	nended Use	ful Reagents			
ANTIGEN	RETRIEVAL BI	UFFER, pH8.0 (BUF025	<u>5A)</u>		
North & South America	Tel: +1 800 265 73 Fax: +1 919 878 3	3751	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
		ales_us@bio-rad.com	Email: antibody_sales_uk@bio		Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specific	> datasheet for this prod	luct, please use our online 'M383344:210513'	search tool at: b	io-rad-antibodies.com/datasheets
			Printed on 01 Mar 2024		

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint