

Datasheet: AHP2116

Description:	GOAT ANTI HUMAN DESMIN (C-TERMINAL)
Specificity:	DESMIN (C-TERMINAL)
Format:	Purified
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal						
	communications from the originators. Please refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.						
	Flow Cytometry			•			
		Immunohistology - Frozen					
	Immunohistology - Paraffin (1)	•			3.0 - 6.0ug/ml		
	ELISA	-			1/2000		
	Immunoprecipitation			-			
	Western Blotting	-			0.1 - 0.3ug/ml		
	 Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) This product requires heat-mediated antigen retrieval prior to staining of paraffin sections. Sodium citrate buffer pH6.0 is recommended for this purpose. 						
Target Species	Human						
Species Cross Reactivity	s Reacts with: Mouse, Rat, Pig Based on sequence similarity, is expected to react with:Dog, Bovine						
-	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for						
Product Form	further information. Purified IgG - liquid						

Antiserum Preparation Antiserum to human Desmin (CT) was raised by repeated immunisation of goats with

	highly purified antigen. Purified IgG was prepared by affinity chromatography.
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Synthetic peptide sequence C-RDGEVVSEATQQQHE from the C-Terminal region of Desmin (NP_001918.3).
External Database Links	UniProt: P17661 Related reagents Entrez Gene: 1674 DES Related reagents
RRID	AB_2092438
Specificity	 Goat anti Human Desmin antibody recognizes an epitope within the C-terminal (CT) region of human Desmin, a type III intermediate filament (IF), localized near the Z line in sacromeres of skeletal, smooth and cardiac muscle tissue. Desmin plays an essential role in the architecture of muscle cells, connecting the Z-disk to the subsarcolemmal cytoskeleton, and the contractile apparatus to the, mitochondria, cell
	nucleus and post-synaptic areas of motor endplates. Desmin is an early indicator for muscle tissue in embryogenesis, and a marker for tumours of myogenic origin. Mutations in the Desmin gene, are responsible for the neuromuscular disorder known as desmin- related cardio skeletal myopathy (CSM), characterized by skeletal muscle weakness.
Histology Positive Control Tissue	Human prostate
Western Blotting	Goat anti Human Desmin detects a band of approximately 55kDa in human skeletal muscle cell lysates.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10058 available at:

Informati	<u> </u>	ttps://www.bio 0058	<u>-rad-antibod</u>	ies.com/SDS/AF	<u>IP2116</u>	
Regulato	ry F	or research p	urposes only			
Relate	d Products					
Recomm	nended Seco	ndary Antib	odies			
Rabbit A	nti Goat IgG (Fo	c) (STAR122	.) <u>FITC</u> , <u>HRP</u>	-		
Recomm	nended Usefi	ul Reagents				
	RETRIEVAL BU	<u>FFER, pH8.0 (E</u>	<u>3UF025A)</u>			
MATIOLIN						
	Tel: +1 800 265 737 Fax: +1 919 878 375			+44 (0)1865 852 700 +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint