

Datasheet: AHP2167T

BATCH NUMBER 1608

Description:	RABBIT ANTI HUMAN MAP1LC3A/B (N-TERMINAL)
Specificity:	MAP1LC3A/B (N-TERMINAL)
Other names:	Atg8-LC3
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1/1000 - 1/5000
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Mouse
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form Purified IgG - liquid

Antiserum Preparation Antiserum to human LC3A was raised by repeated immunisation of rabbits with highly

purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
---------------------------------	--

Approx. Protein Concentrations	IgG concentration 1.0mg/ml
---------------------------------------	----------------------------

Immunogen	Synthetic peptide sequence PSDRPFKQRRSFADC from the N-Terminal region of LC3A (NP_115903.1; NP_852610.1).
------------------	---

External Database Links	UniProt: Q9H492 Related reagents Q9GZQ8 Related reagents Q91VR7 Related reagents Q9CQV6 Related reagents Entrez Gene: 84557 MAP1LC3A Related reagents 81631 MAP1LC3B Related reagents 66734 Map1lc3a Related reagents 67443 Map1lc3b Related reagents
--------------------------------	--

Synonyms	Map1alc3, MAP1ALC3, Map1lc3
-----------------	-----------------------------

RRID	AB_10673306
-------------	-------------

Specificity	<p>Rabbit anti Human MAP1LC3A/B (N-Terminal) antibody specifically recognizes an epitope within the N-Terminal (NT) region of both MAP1LC3A (Microtubule-associated proteins 1A/1B light chain 3A/LC3A) and MAP1LC3B (Microtubule-associated proteins 1A/1B light chain 3B/LC3B), ubiquitin-like proteins and members of the MAP1LC3 family, which are widely used as reliable markers for the monitoring of autophagy.</p> <p>LC3-I is the cytosolic form of LC3, which is converted into the active, membrane-bound form LC3-II, during the autophagy process. Tracking the level of conversion of LC3-I to LC3-II provides an indicator of autophagic activity, and levels of LC3-II in particular, correlate with the extent of autophagosome formation, due to its association with the autophagosome membrane.</p> <p>Rabbit anti Human MAP1LC3A/B (N-Terminal) antibody recognizes both the LC3-I and LC3-II forms of MAP1LC3A and MAP1LC3B.</p>
--------------------	--

Western Blotting	AHP2167T detects a band of approximately 14-15kDa corresponding to LC3-II, and a band of approximately 17kDa corresponding to LC3-I, in HeLa cell lysates.
-------------------------	--

References

1. Iwata, A. *et al.* (2005) HDAC6 and microtubules are required for autophagic degradation of aggregated huntingtin. [J Biol Chem. 280 \(48\): 40282-92.](#)
2. Riley, B.E. *et al.* (2010) Ubiquitin accumulation in autophagy-deficient mice is dependent on the Nrf2-mediated stress response pathway: a potential role for protein aggregation in autophagic substrate selection. [J Cell Biol. 191 \(3\): 537-52.](#)
3. Gjyshi, O. *et al.* (2015) Kaposi's Sarcoma-Associated Herpesvirus Induces Nrf2 Activation in Latently Infected Endothelial Cells through SQSTM1 Phosphorylation and Interaction with Polyubiquitinated Keap1. [J Virol. 89: 2268-86](#)
4. Huang, L. *et al.* (2014) AKI after conditional and kidney-specific knockdown of stanniocalcin-1. [J Am Soc Nephrol. 25: 2303-15.](#)
5. Girard, B.J. *et al.* (2015) Cytoplasmic PELP1 and ERRgamma Protect Human Mammary Epithelial Cells from Tam-Induced Cell Death. [PLoS One. 10 \(3\): e0121206.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/AHP2167T>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M373279:200907'

Printed on 17 Jan 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)