

Datasheet: AHP2167T

Description:	RABBIT ANTI HUMAN MAP1LC3A/B (N-TERMINAL)
Specificity:	MAP1LC3A/B (N-TERMINAL)
Other names:	Atg8-LC3
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1/1000 - 1/5000
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Mouse
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form Purified IgG - liquid

Antiserum Preparation Antiserum to human LC3A was raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Synthetic peptide sequence PSDRPFKQRRSFADC from the N-Terminal region of LC3A (NP_115903.1; NP_852610.1).
External Database Links	<p>UniProt:</p> <p>Q9H492 Related reagents</p> <p>Q9GZQ8 Related reagents</p> <p>Q91VR7 Related reagents</p> <p>Q9CQV6 Related reagents</p> <p>Entrez Gene:</p> <p>84557 MAP1LC3A Related reagents</p> <p>81631 MAP1LC3B Related reagents</p> <p>66734 Map1lc3a Related reagents</p> <p>67443 Map1lc3b Related reagents</p>
Synonyms	Map1alc3, MAP1ALC3, Map1lc3
RRID	AB_10673306
Specificity	<p>Rabbit anti Human MAP1LC3A/B (N-Terminal) antibody specifically recognizes an epitope within the N-Terminal (NT) region of both MAP1LC3A (Microtubule-associated proteins 1A/1B light chain 3A/LC3A) and MAP1LC3B (Microtubule-associated proteins 1A/1B light chain 3B/LC3B), ubiquitin-like proteins and members of the MAP1LC3 family, which are widely used as reliable markers for the monitoring of autophagy.</p> <p>LC3-I is the cytosolic form of LC3, which is converted into the active, membrane-bound form LC3-II, during the autophagy process. Tracking the level of conversion of LC3-I to LC3-II provides an indicator of autophagic activity, and levels of LC3-II in particular, correlate with the extent of autophagosome formation, due to its association with the autophagosome membrane.</p> <p>Rabbit anti Human MAP1LC3A/B (N-Terminal) antibody recognizes both the LC3-I and LC3-II forms of MAP1LC3A and MAP1LC3B.</p>
Western Blotting	AHP2167T detects a band of approximately 14-15kDa corresponding to LC3-II, and a band of approximately 17kDa corresponding to LC3-I, in HeLa cell lysates.
References	1. Iwata, A. <i>et al.</i> (2005) HDAC6 and microtubules are required for autophagic degradation of aggregated huntingtin. J Biol Chem. 280 (48): 40282-92.

2. Riley, B.E. *et al.* (2010) Ubiquitin accumulation in autophagy-deficient mice is dependent on the Nrf2-mediated stress response pathway: a potential role for protein aggregation in autophagic substrate selection. [J Cell Biol. 191 \(3\): 537-52.](#)
3. Gjyshi, O. *et al.* (2015) Kaposi's Sarcoma-Associated Herpesvirus Induces Nrf2 Activation in Latently Infected Endothelial Cells through SQSTM1 Phosphorylation and Interaction with Polyubiquitinated Keap1. [J Virol. 89: 2268-86](#)
4. Huang, L. *et al.* (2014) AKI after conditional and kidney-specific knockdown of stanniocalcin-1. [J Am Soc Nephrol. 25: 2303-15.](#)
5. Girard, B.J. *et al.* (2015) Cytoplasmic PELP1 and ERRgamma Protect Human Mammary Epithelial Cells from Tam-Induced Cell Death. [PLoS One. 10 \(3\): e0121206.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
- Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
- Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
- Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
- Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@680](#), [DyLight@800](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M382552:210513'

Printed on 18 May 2021