

Datasheet: AHP2268 BATCH NUMBER 158644

| Description: | GOAT ANTI WNT-3 |
|---------------|---------------------|
| Specificity: | WNT-3 |
| Other names: | INT4 |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | | |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin (1) | • | | | 3.0 - 5.0ug/ml |
| ELISA | • | | | 1:16000 |
| Immunoprecipitation | | | | |
| Western Blotting | • | | | 1.0 - 3.0ug/ml |
| Immunofluorescence | • | | | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species | Human |
|-----------------------------|---|
| Species Cross Reactivity | Reacts with: Mouse Based on sequence similarity, is expected to react with:Dog, Rat |
| | N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for |
| | further information. |

| Product Form | Purified IgG - liquid |
|-----------------------|--|
| Antiserum Preparation | Antiserum to Wnt-3 antigen. Purified Ig0 peptide |

Antiserum to Wnt-3 was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography using the immunizing

Preservative Stabilisers

Buffer Solution

0.02% Sodium Azide (NaN₃) 0.5% Bovine Serum Albumin

TRIS buffered saline

Approx. Protein Concentrations

IgG concentration 0.5 mg/ml

Immunogen

Synthetic peptide sequence C-GRGHNTRTEKRKEK from the internal region of Wnt-3 (NP 110380.1).

External Database Links

UniProt:

P56703 Related reagents

Entrez Gene:

7473 WNT3 Related reagents

Synonyms

INT4

Specificity

Goat anti Wnt-3 antibody recognises both human and mouse proto-oncogene Wnt-3, a member of the Wnt family, and ligand of the frizzled family of seven transmembrane receptors.

Wnt-3, otherwise known as Int-4, is involved in cell fate decisions and patterning during embryogenesis, cell signalling during neural tube development, and implicated in the tumorogenesis of several cancers including breast, lung, hepatocellular(HCC), and gastric carcinoma.

Over-expression of Wnt-3 playes a pivotal role in the resistance of HER2-overexpressing breast tumors to the breast cancer treatment drug Trastuzumab, also known as Herceptin (<u>Wu et al. 2012</u>). The effects of Wnt-3 over-expression included the transactivated expression of EGFR (epidermal growth factor receptor) an increased expression of B-catenin within the nucleus, an increase in N-cadherin and the neural crest transcription factor SLUG, as well as promoting an EMT (epithelial to mesenchymal transition) response.

Goat anti Wnt-3 polyclonal antibody exhibits strong fibrous staining in myoepithelial cells of human breast tissue in paraffin sections.

Goat anti Wnt-3 polyclonal antibody stains mouse hippocampal primary astrocytes in ICC/IF (Okamoto et al. 2011).

| Histology Positive Control Tissue | Human breast | |
|--------------------------------------|--|--------------|
| Western Blotting | AHP2268 detects a band of approximately 40kDa in HEK293 | cell lysates |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. | |
| Guarantee | 12 months from date of despatch | |
| Health And Safety Information | Material Safety Datasheet documentation #10058 available at https://www.bio-rad-antibodies.com/SDS/AHP2268 10058 | : |
| Regulatory | For research purposes only | |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382847:210513'

Printed on 21 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint