

Datasheet: AHP2311

Description:	RABBIT ANTI HISTONE DEACETYLASE 4
Specificity:	HISTONE DEACETYLASE 4
Other names:	HDAC4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	•			5 ug/ml
Western Blotting	•			0.5 - 4.0 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Species Cross Reactivity	Reacts with: Mouse Based on sequence similarity, is expected to react with:Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species.	
Product Form	Purified IgG - liquid	
Buffer Solution	Phosphate buffered saline	
Preservative	50% Glycerol	
Stabilisers	1% Bovine serum Albumin	
	0.02% sodium azide	
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml	
Immunogen	Synthetic peptide corresponding to residues surrounding amino acid 10 of human HDAC4	
External Database Links	UniProt:	

Related reagents

**Entrez Gene:** 

P56524

## 9759 HDAC4 Related reagents

Synonyms	KIAA0288					
Specificity	Rabbit anti histone deacetylase 4 antibody recognizes histone deacetylase 4, also known as HDAC4. Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by HDAC4 belongs to the class II histone deacetylase/acuc/apha family and has been implicated to play a role in muscle maturation.					
Western Blotting	Rabbit anti histone deacetylase 4 recognizes a band of approximately 140 kDa in mouse brain tissue lysate					
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.					
Guarantee	12 months from date of despatch					
Health And Safety Information	Material Safety Datasheet documentation #10048 available at:  10048: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</a>					
Regulatory	For research purposes only					

# **Related Products**

# **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) **FITC** Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) **RPE** 

Sheep Anti Rabbit IgG (STAR36...) DyLight®488, DyLight®680, DyLight®800

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M361318:200210'

Printed on 01 May 2020

© 2020 Bio-Rad Laboratories Inc | Legal | Imprint