

Datasheet: AHP2312

Description:	RABBIT ANTI HISTONE DEACETYLASE 4
Specificity:	HISTONE DEACETYLASE 4
Other names:	HDAC4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	▪			5 ug/ml
Western Blotting	▪			0.5 - 4.0 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Based on sequence similarity, is expected to react with:Chicken N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Buffer Solution	Phosphate buffered saline
Preservative	30% Glycerol
Stabilisers	0.5% Bovine Serum Albumin 0.01% Thiomersal
Approx. Protein Concentrations	IgG concentration 0.2 mg/ml
Immunogen	Synthetic peptide surrounding amino acid 10 of human HDAC4
External Database Links	UniProt: P56524 Related reagents Entrez Gene:

Synonyms	KIAA0288
-----------------	----------

Specificity	Rabbit anti histone deacetylase 4 antibody recognizes histone deacetylase 4, also known as HDAC4. Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by HDAC4 belongs to the class II histone deacetylase/acuc/apha family and has been implicated to play a role in muscle maturation.
--------------------	--

Western Blotting	Rabbit anti histone deacetylase 4 recognizes a band of approximately 140 kDa in Jurkat cell lysate
-------------------------	--

Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10096 available at: 10096: https://www.bio-rad-antibodies.com/uploads/MSDS/10096.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@680](#), [DyLight@800](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M361319:200210'

Printed on 01 May 2020