

Datasheet: AHP2366

Description:	RABBIT ANTI BOVINE MCP-1
Specificity:	MCP-1
Other names:	CCL2
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.5 ug/ml - 10 ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 ug/ml - 1.0 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	Reacts with: Horse, Pig N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Antiserum Preparation	Antisera to bovine MCP-1 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.8 mg/ml
Immunogen	Recombinant bovine MCP-1

**External Database
Links**

UniProt:

[P28291](#) [Related reagents](#)

Entrez Gene:

[281043](#) CCL2 [Related reagents](#)

Synonyms

SCYA2

Specificity

Rabbit anti Bovine MCP-1 antibody recognizes bovine monocyte chemotactic protein-1 (MCP-1), also known as C-C motif chemokine 2 (CCL2). Bovine MCP-1 is a 99 amino acid, 11.1 kDa secreted protein.

The chemokine MCP-1 is expressed by a range of cells including monocytes, vascular endothelial cells, smooth muscle cells, glomerular mesangial cells and osteoblasts.

MCP-1 is thought to play an important role in the inflammatory response of blood monocytes and tissue macrophages.

ELISA

This product may be used as a capture antibody in a sandwich ELISA together with [AHP2367B](#) as the detection reagent. Recombinant bovine MCP-1 ([PBP024](#)) may be used as a standard.

Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@680](#), [DyLight@800](#)

Recommended Useful Reagents

[RECOMBINANT BOVINE MCP-1 \(PBP024\)](#)

[GOAT ANTI BOVINE MCP-1:Biotin \(AHP2367B\)](#)

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M334152:181127'

