

Datasheet: AHP2395B BATCH NUMBER 147890

Description:	RABBIT ANTI PIG INTERLEUKIN-17A:Biotin
Specificity:	IL-17A
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA				0.05ug/ml - 0.5ug/ml
Immunoprecipitation			•	
Western Blotting			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig	
Species Cross Reactivity	Reacts with: Horse, Sheep, Bovine, Human, Rat N.B. Antibody reactivity and working conditions may vary betwee reactivity is derived from testing within our laboratories, peer-respersional communications from the originators. Please refer to refurther information.	viewed publications or
Product Form	Purified IgG conjugated to Biotin - liquid	
Antiserum Preparation	Antisera to pig interleukin-17A were raised by repeated immunis purified antigen. Purified IgG was prepared from whole serum b	= -
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant pig interleukin-17A
External Database Links	UniProt:
LITIKS	Q60I29 Related reagents
	Entrez Gene:
	449530 IL17A Related reagents
Synonyms	IL17
Specificity	Rabbit anti Pig interleukin-17A antibody recognizes porcine interleukin-17A (IL-17A), a 35 kDa disulfide-linked homodimer consisting of two 153 amino acid chains, which belongs to the IL-17 family (classified as IL-17A to IL-17F), and signals through the IL-17
	receptor (IL-17R/CDw217).
	receptor (IL-17R/CDw217). IL-17, originally thought to have been produced primarily by activated CD4+ T cells, appears to have a range of cellular origins and induces a variety of target cells to secrete inflammatory cytokines, including IL-1β, IL-6, TNFα, IFNγ and granulocyte colonystimulating factor, as well as synergizing with other inflammatory cytokines and agonists.
ELISA	IL-17, originally thought to have been produced primarily by activated CD4+ T cells, appears to have a range of cellular origins and induces a variety of target cells to secrete inflammatory cytokines, including IL-1 β , IL-6, TNF α , IFN γ and granulocyte colony-
ELISA Storage	IL-17, originally thought to have been produced primarily by activated CD4+ T cells, appears to have a range of cellular origins and induces a variety of target cells to secrete inflammatory cytokines, including IL-1β, IL-6, TNFα, IFNγ and granulocyte colonystimulating factor, as well as synergizing with other inflammatory cytokines and agonists. This product may be used as a detection antibody in a sandwich ELISA together with AHP2395 as the capture reagent. Recombinant pig interleukin-17A (PPP028) may be
Storage	IL-17, originally thought to have been produced primarily by activated CD4+ T cells, appears to have a range of cellular origins and induces a variety of target cells to secrete inflammatory cytokines, including IL-1β, IL-6, TNFα, IFNγ and granulocyte colonystimulating factor, as well as synergizing with other inflammatory cytokines and agonists. This product may be used as a detection antibody in a sandwich ELISA together with AHP2395 as the capture reagent. Recombinant pig interleukin-17A (PPP028) may be used as a standard. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we
	IL-17, originally thought to have been produced primarily by activated CD4+ T cells, appears to have a range of cellular origins and induces a variety of target cells to secrete inflammatory cytokines, including IL-1β, IL-6, TNFα, IFNγ and granulocyte colonystimulating factor, as well as synergizing with other inflammatory cytokines and agonists. This product may be used as a detection antibody in a sandwich ELISA together with AHP2395 as the capture reagent. Recombinant pig interleukin-17A (PPP028) may be used as a standard. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

North & South Tel: +1 800 265 7376
America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M334205:181127'

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint