

# Datasheet: AHP2561 BATCH NUMBER 154700

Description:	RABBIT ANTI BRCA1 (pSer988)
Specificity:	BRCA1 (pSer988)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μΙ

## **Product Details**

**Applications** 

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/500 - 1/2000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid

Antiserum Preparation Antiserum to BRCA1 (pSer988) was raised by repeated immunization of rabbits with

highly purified antigen. Purified IgG was prepared from whole serum by affinity

chromatography.

 Buffer Solution
 Phosphate buffered saline

 Preservative
 0.02% Sodium Azide

 Stabilisers
 50% Glycerol

Immunogen Phospho specific-peptide corresponding to residues surrounding serine 988 of human

BRCA1

**External Database** 

Links UniProt:

P38398 Related reagents

#### **Entrez Gene:**

672 BRCA1 Related reagents

Rabbit anti BRCA1 (pSer988) antibody recognizes BRCA1, also known as breast cancer type 1 susceptibility protein and RING finger protein 53, when phosphorylated at serine 988. Breast cancer susceptibility genes BRCA1 and BRCA2 are associated with hereditary breast and ovarian cancers. BRCA1 is a tumor suppressor gene encoding an E3 ubiquitin protein ligase that plays a major role in DNA damage repair (Roy et al. 2011)  BRCA1 is also involved in the regulation of multiple processes including transcription and cell cycle progression. BRCA1 co-localizes with Chk2 in nuclear foci, in response to DNA damage BCRA1 is phosphorylated at serine 988 by Chk2. This phosphorylation event results in the release and dispersion of BRCA1 and is suggested to be important for the function of BRCA1 in the DNA damage response (Lee et al. 2000).  Further Reading  1. Roy, R. et al. (2011) BRCA1 and BRCA2: different roles in a common pathway of genome protection. Nat Rev Cancer. 12 (1): 68-78. 2. Lee, J.S. et al. (2000) hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. Nature. 404 (6774): 201-4.  Storage  Storage Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  Guarantee  12 months from date of despatch  Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/AHP2561 10049		
cancer type 1 susceptibility protein and RING finger protein 53, when phosphorylated at serine 988. Breast cancer susceptibility genes BRCA1 and BRCA2 are associated with hereditary breast and ovarian cancers. BRCA1 is a tumor suppressor gene encoding an E3 ubiquitin protein ligase that plays a major role in DNA damage repair (Roy et al. 2011)  BRCA1 is also involved in the regulation of multiple processes including transcription and cell cycle progression. BRCA1 co-localizes with Chk2 in nuclear foci, in response to DNA damage BCRA1 is phosphorylated at serine 988 by Chk2. This phosphorylation event results in the release and dispersion of BRCA1 and is suggested to be important for the function of BRCA1 in the DNA damage response (Lee et al. 2000).  Further Reading  1. Roy, R. et al. (2011) BRCA1 and BRCA2: different roles in a common pathway of genome protection. Nat Rev Cancer. 12 (1): 68-78. 2. Lee, J.S. et al. (2000) hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. Nature, 404 (6774): 201-4.  Storage  Store at -20°C  Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  12 months from date of despatch  Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/AHP2561 10049	Synonyms	RNF53
cell cycle progression. BRCA1 co-localizes with Chk2 in nuclear foci, in response to DNA damage BCRA1 is phosphorylated at serine 988 by Chk2. This phosphorylation event results in the release and dispersion of BRCA1 and is suggested to be important for the function of BRCA1 in the DNA damage response (Lee et al. 2000).  Further Reading  1. Roy, R. et al. (2011) BRCA1 and BRCA2: different roles in a common pathway of genome protection. Nat Rev Cancer. 12 (1): 68-78. 2. Lee, J.S. et al. (2000) hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. Nature. 404 (6774): 201-4.  Storage  Storage Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  Guarantee  12 months from date of despatch  Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/AHP2561 10049	Specificity	cancer type 1 susceptibility protein and RING finger protein 53, when phosphorylated at serine 988. Breast cancer susceptibility genes BRCA1 and BRCA2 are associated with
genome protection. Nat Rev Cancer. 12 (1): 68-78.  2. Lee, J.S. et al. (2000) hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. Nature. 404 (6774): 201-4.  Storage  Store at -20°C  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  Guarantee  12 months from date of despatch  Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/AHP2561 10049		results in the release and dispersion of BRCA1 and is suggested to be important for the
Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  12 months from date of despatch  Health And Safety Information  Material Safety Datasheet documentation #10049 available at:  https://www.bio-rad-antibodies.com/SDS/AHP2561 10049	Further Reading	genome protection. Nat Rev Cancer. 12 (1): 68-78.  2. Lee, J.S. et al. (2000) hCds1-mediated phosphorylation of BRCA1 regulates the DNA
Health And Safety Material Safety Datasheet documentation #10049 available at:  https://www.bio-rad-antibodies.com/SDS/AHP2561  10049	Storage	Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
https://www.bio-rad-antibodies.com/SDS/AHP2561 10049	Guarantee	12 months from date of despatch
Regulatory For research purposes only	Health And Safety Information	https://www.bio-rad-antibodies.com/SDS/AHP2561
	Regulatory	For research purposes only

## **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR124...) <u>HRP</u> Sheep Anti Rabbit IgG (STAR35...) <u>RPE</u>

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

**Recommended Useful Reagents** 

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

'M361331:200210'

Email: antibody\_sales\_de@bio-rad.com

### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint