

Datasheet: AHP2585

Description:	RABBIT ANTI E3 UBIQUITIN-PROTEIN LIGASE CBL (pTyr700)
Specificity:	E3 UBIQUITIN-PROTEIN LIGASE CBL (pTyr700)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 μl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin				1/50 - 1/100
Western Blotting	•			1/500 - 1/1000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid

Antiserum Preparation Antiserum to E3 ubiqutin-protein ligase CBL (pTyr700) was raised by repeated

immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole

serum by affinity chromatography.

UniProt: Links

> P22681 Related reagents

Entrez Gene:

867 CBL Related reagents

Synonyms	CBL2, RNF55
Specificity	Rabbit anti E3 ubiquitin-protein ligase CBL (pTyr700) antibody recognizes the Casitas B-lineage lymphoma (CBL) E3 ubiquitin ligase, also known as c-CBL proto-oncogene and ring finger protein 55, when phosphorylated at tyrosine 700. The CBL protein is a RING finger E3-ubiquitin ligase that plays a key role in signal transduction through mediating monoubiquitination and endocytosis of receptor tyrosine kinases (Haglund et al. 2002, Dikic and Giordano 2003).
	E3 ubiqutin-protein ligase CBL is phosphorylated at tyrosine 700 by Fyn, Yes and Syk (Feshchenko et al. 1998).
Further Reading	 Haglund, K. <i>et al.</i> (2002) Cbl-directed monoubiquitination of CIN85 is involved in regulation of ligand-induced degradation of EGF receptors. Proc Natl Acad Sci U S A. 99 (19): 12191-6. Dikic, I. & Giordano, S. (2003) Negative receptor signalling. Curr Opin Cell Biol. 15 (2): 128-35. Feshchenko, E.A. <i>et al.</i> (1998) Fyn, Yes, and Syk phosphorylation sites in c-Cbl map to the same tyrosine residues that become phosphorylated in activated T cells. J Biol Chem. 273 (14): 8323-31.
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) FITC

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Sheep Anti Rabbit IgG (STAR36...) <u>DyLight®488</u>, <u>DyLight®680</u>, <u>DyLight®800</u>

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M354072:190624'

Printed on 29 Aug 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint