

Datasheet: AHP2700

Description:	RABBIT ANTI SMAD2 (pSer465/pSer467)/SMAD3 (pSer423/pSer425)
Specificity:	SMAD2 (pSer465/pSer467)/SMAD3 (pSer423/pSer425)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/500 - 1/2000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG - liquid

Antiserum Preparation Antiserum to SMAD2 (pSer465/pSer467)/SMAD3 (pSer423/pSer425) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution Phosphate buffered saline

Preservative 0.02% Sodium Azide
Stabilisers 50% Glycerol

Immunogen Phospho specific-peptide corresponding to residues surrounding serine 465/467 of human SMAD2 and serine 423/425 of human SMAD3

External Database Links

UniProt:

[Q15796](#) [Related reagents](#)

[P84022](#) [Related reagents](#)

Entrez Gene:

[4087](#) SMAD2 [Related reagents](#)

[4088](#) SMAD3 [Related reagents](#)

Synonyms	MADH2, MADH3, MADR2
Specificity	Rabbit anti SMAD2 (pSer465/pSer467)/SMAD3 (pSer423/pSer425) antibody recognizes mothers against decapentaplegic homolog 2 (SMAD2), when phosphorylated at serine 465 and serine 467 and mothers against decapentaplegic homolog 3 (SMAD3) when phosphorylated at serine 423 and serine 425. The SMAD protein family mediates transforming growth factor-beta (TGF- β) signaling from the membrane to the nucleus (Attisano, L. & Wrana, J.L. 2002). SMAD2 is phosphorylated on serine 465 and serine 467 by TGF beta type I receptor (TbetaRI). This phosphorylation is essential for SMAD2's binding to SMAD4 and for the orchestration of TGF beta signaling (Abdollah, S. et al. 1997). The antibody also recognizes SMAD3 phosphorylated on Ser423 and Ser425 by homology. The TGF beta signaling pathway is involved in many cellular processes including cell growth, differentiation and proliferation. The pathway is deregulated in various diseases, including cancer.
Storage	Store at -20°C Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: 10049: https://www.bio-rad-antibodies.com/uploads/MSDS/10049.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[RABBIT ANTI SMAD2 \(VPA00182\)](#)

[RABBIT ANTI SMAD3 \(VPA00166\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M340143:190104'

Printed on 01 May 2020