

Datasheet: AHP3011

Description:	on: GOAT ANTI SCRIB	
Specificity:	SCRIB	
Format:	Purified	
Product Type:	Polyclonal Antibody	
Isotype:	Polyclonal IgG	
Quantity:	0.1 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			3 μg/ml - 0.5 μg/ml
Immunofluorescence	•			10 μg/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid

Antiserum Preparation Antisera to human SCRIB were raised by repeated immunizations of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.02% Sodium Azide 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Peptide with sequence C-PEGPGKEKELPGQ, from the internal region of the protein sequence according to NP_874365.3; NP_056171.3.
Specificity	Goat anti Human SCRIB antibody recognizes protein scribble homolog, also known as

Scribble, CRIB1, KIAA0147, LAP4, SCRB1, and VARTUL.

SCRIB is a scaffold protein responsible for correct apical-basal cell polarity of the epithelial tissues (Kapil et al.. 2017). Appropriate polarity is important for cell migration and morphogenesis during development, and for wound healing in adult tissue. SCRIB also plays an important role in preventing cancer progression. Loss of SCRIB function disrupts apical-basal polarity and junctional integrity, and causes inappropriate proliferation which results in overgrowth of tissues. Overexpression of SCRIB has been found in tumors of the colon, breast, lung, ovary and prostate (Ono et al.. 2015). Correct SCRIB function depends on its localization to the plasma membrane, with mislocalization promoting invasion of cells through the extracellular matrix (Sun et al., 2017). Accordingly, SCRIB can be found in the cytoplasm in some tumor types, and even in the nucleus in hepatocellular carcinoma and liver tumors (Kapil et al., 2017). In Drosophila, SCRIB has been found to orchestrate an intracellular signaling process for forgetting information (Cervantes-Sandoval et al.. 2016).

Western Blotting

Goat anti Human SCRIB antibody detects a band of approximately 200 to 250 kDa in cell

lysates.

Storage

This product is shipped at ambient temperature.

Store at -20°C only.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10058 available at:

https://www.bio-rad-antibodies.com/SDS/AHP3011

10058

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

America

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M439210:250523'

Printed on 23 May 2025