

Datasheet: AHP3049 BATCH NUMBER 160737

Description:	RABBIT ANTI HUMAN HISTONE H3 (Ac)
Specificity:	HISTONE H3 (Ac)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Western Blotting	-			0.2 μg/ml - 1.0 μg/ml		
	Immunofluorescence				1/500		
	Chromatin Immunoprecipitation	-			10 µl per ChIP		
	Where this product has necessarily exclude its u a guide only. It is recomi system using appropriat	use in such mended th	n procedur at the use	es. Suggested workin r titrates the product f	g dilutions are given as		
Target Species	Human						
Product Form	Purified IgG - liquid						
Antiserum Preparatio	on Antiserum to human hist highly purified antigen. F chromatography on Prot	Purified IgC	,				
Buffer Solution	Phosphate buffered sali	ne					
Preservative Stabilisers	0.035% Sodium Azide (I 30% Glycerol	NaN ₃)					
Carrier Free	Yes						
Immunogen	A peptide including acet	ylated lysi	nes contai	ned in the N-terminal	tail of human histone H3		

External Database Links	UniProt: <u>P68431</u> <u>Related reagents</u> Entrez Gene:					
	8968 HIST1H3F Related reagents					
Synonyms	H3FA, H3FB, H3FC, H3FD, H3FF, H3FH, H3FI, H3FJ, H3FK, H3FL					
Specificity	Rabbit anti Human histone H3 (Ac) antibody recognizes histone H3 when acetylated.					
	Histone H3 is one of the four core histones that make up the nucleosome core particle. Nucleosomes are the smallest subunit of chromatin and are made up of 146 bp of DNA wrapped around an octamer comprised of pairs of the four core histones (H2A, H2B, H3, and H4) (Smith, 1991). Lysine residues within the N-terminal tail protruding from the histone core of the nucleosome are acetylated and deacetylated as part of gene regulation. Hyperacetylation of histone tails makes chromatin more accessible to DNA-binding proteins by weakening histone-DNA and nucleosome-nucleosome interactions (Kuo and Allis, 1998).					
	Wide species cross-reactivity is expected from Rabbit anti Human histone H3 (Ac) antibody based on sequence.					
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.					
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.					
Guarantee	12 months from date of despatch					
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/AHP3049 10049					
Regulatory	For research purposes only					

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34)	<u>FITC</u>			
Sheep Anti Rabbit IgG (STAR35)	<u>RPE</u>			
Goat Anti Rabbit IgG (H/L) (STAR124) <u>HRP</u>				
Goat Anti Rabbit IgG (Fc) (STAR121)	Biotin, FITC, HRP			

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21	То		
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	find a		
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com			
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets								
'M394880:220218'								

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint